

MLX Antibody

Mouse Monoclonal Antibody [Clone PCRP-MLX-1G8]



Catalog No	Format	Size
6945-MSM1-P0	Purified Ab with BSA and Azide	200ug/ml
6945-MSM1-P1	Purified Ab with BSA and Azide	200ug/ml
6945-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Flow Cytometry (Flow)	1-2ug/million cells
Immunofluorescence (IF)	1-3ug/ml
Western Blot (WB)	2-4ug/ml

Product Details	
Clone	PCRP-MLX-1G8
Gene Name	MLX
Immunogen	Recombinant full-length human MLX protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b
Mol. Weight of Antigen	30kDa
Cellular Localization	Cytoplasm, Nucleus
Species Reactivity	Human
Positive Control	HeLa or MCF7 cells.


**Optimal dilution for a specific application should be determined.*

Product Images for MLX Antibody



Immunofluorescence Analysis of PFA-fixed HeLa cells. MLX Mouse Monoclonal Antibody (PCRP-MLX-1G8) followed by goat anti-mouse IgG-CF488 (green); counterstain phalloidin (red).

SDS-PAGE Analysis of Purified MLX Mouse Monoclonal Antibody (PCRP-MLX-1G8). Confirmation of Purity and Integrity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteins using MLX Mouse Monoclonal Antibody (PCRP-MLX-1G8). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Flow Cytometric Analysis of PFA-fixed HeLa cells. MLX Mouse Monoclonal Antibody (PCRP-MLX-1G8) followed by goat anti-mouse IgG-CF488 (blue); unstained cells (red).

Specificity & Comments

Max is a nuclear localized bHLH-Zip protein that forms homodimers or heterodimers with Myc family members, including Myc, Mad1, Mad3, Mad4, Mxi1 and Mnt (or Rox). These dimers bind to the E-box sequence CACGTG in order to regulate cell growth, proliferation and apoptosis. Mlx (Max-like protein X) is a bHLH-Zip protein that is structurally and functionally related to Max. Like Max, Mlx is broadly expressed in many tissues and has a long half-life. Mlx also forms homodimers or heterodimers with members of the Myc family, specifically Mad1, Mad4 and Rox, and members of the Mondo family, to repress or activate transcription from CACGTG E-boxes. MondoA forms weak homodimers and preferentially forms heterodimers with Mlx. The MondoA/Mlx complex is primarily localized to the cytoplasm, but will translocate to the nucleus in response to leptomycin B. Mlx can also dimerize with WBSCR14, a protein involved in Williams-Beuren syndrome (WBS), to repress E-box transcription, which provides further evidence that Mlx is a critical element in a transcription factor network.

Research Areas

Apoptosis, Autophagy, Nuclear Marker

Known Applications & Suggested Dilutions

Flow Cytometry (1-2ug/million cells) | Immunofluorescence (1-2ug/ml) | Western Blot (1-2ug/ml) | .Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.