

TGF-alpha (Transforming Growth Factor alpha) Antibody

Mouse Monoclonal Antibody [Clone TG86]

Catalog No	Format	Size
7039-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
7039-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
7039-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dilution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	TG86
Gene Name	TGFA
Immunogen	A 10-amino acid synthetic peptide (aa34-43; PPVAAAVVSH) from human TGF?
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	6kDa
Cellular Localization	Cell membrane, Extracellular space, Secreted
Species Reactivity	Human, Rabbit, Zebrafish
Positive Control	Jurkat or Ramos cells. Testicular Carcinoma.

*Optimal dilution for a specific application should be determined.

Product Images for TGF-alpha (Transforming Growth Factor alpha) Antibody



Formalin-paraffin human Testicular Carcinoma Stained with TGF alpha MAb (TG86)

Specificity & Comments

This antibody reacts with the TGF alpha and shows no cross-reaction with EGF and the neuropeptide synenkephalin. The staining with this MAb is completely blocked by the peptide used for raising this antibody. TGF (aa50) is a growth factor with 33% homology to EGF, binds to EGFR, activates tyrosine phosphorylation of the receptor, and stimulates cell proliferation. It plays a role in tumor initiation by inducing the reversible transformed phenotype.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

AKT Signaling, Angiogenesis, Cardiovascular, Infectious Disease, Signal Transduction, Transcription Factors

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
