

TIMP2 (Tissue Inhibitor of Metalloproteinase 2) Antibody

Mouse Monoclonal Antibody [Clone SPM356]

Catalog No	Format	Size
7077-MSM1X-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
7077-MSM1X-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
7077-MSM1X-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	SPM356
Gene Name	TIMP2
Immunogen	A synthetic peptide from the N-terminus of human TIMP2 protein (aa56-66) (DSGNDIYGNPI)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	21kDa
Cellular Localization	Secreted
Species Reactivity	Cow, Guinea Pig, Human, Mouse, Rabbit, Rat, Zebrafish
Positive Control	Colon, Endometrial, HeLa cells. Normal Placenta. Breast, Prostate or Ovarian Carcinoma.

*Optimal dilution for a specific application should be determined.

Product Images for TIMP2 (Tissue Inhibitor of Metalloproteinase 2) Antibody



Formalin-fixed, paraffin-embedded human Placenta stained with TIMP2 Monoclonal Antibody (SPM356).

Specificity & Comments

It recognizes a protein of 21kDa, identified as tissue inhibitor of metalloproteinases-2 (TIMP-2). It is closely related to TIMP-1 and shows the highest binding affinity to both the latent (pro) and active forms of 72kDa Type IV collagenase (also known as MMP-2 or gelatinase A). It also has affinity for the active form of 92kDa Type IV collagenase (also known as MMP-9 or gelatinase B). TIMP-2 inhibits the proteolytic invasiveness of tumor cells and normal placental trophoblast cells.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

AKT Signaling, Angiogenesis, Cardiovascular, Immunology
