



## TIMP-3 (Tissue Inhibitor of Metalloproteinase-3) Antibody

Catalog No	Format	Size
7078-RBP1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
7078-RBP1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
7078-RBP1-P1ABX	Purified Ab WITHOUT BSA at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details

<b>Gene Name</b>	TIMP3
<b>Immunogen</b>	The amino acid sequence (aa175-211), used as the immunogen for TIMP3 Ab, is 100% homologous in human, cow, dog and horse, and 94% homologous in Mouse and rat.
<b>Host</b>	Rabbit
<b>Clonality</b>	Polyclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	30kDa
<b>Cellular Localization</b>	Extracellular matrix, Extracellular space, Secreted
<b>Species Reactivity</b>	Cow, Dog, Horse, Human, Mouse, Rat
<b>Positive Control</b>	Placenta or breast carcinoma.

*\*Optimal dilution for a specific application should be determined.*

### Product Images for TIMP-3 (Tissue Inhibitor of Metalloproteinase-3) Antibody



Formalin-fixed, paraffin-embedded human placenta stained with TIMP3 Rabbit PAb.

### Specificity & Comments

TIMP3 (tissue inhibitor of metalloproteinases 3), along with family members TIMP1, TIMP2, and TIMP4, are inhibitors of the matrix metalloproteinases (MMPs), a group of peptidases involved in degradation of the extracellular matrix (ECM). An imbalance between MMPs and the associated TIMPs may play a significant role in the invasive phenotype of malignant tumors. TIMP's inhibit the proteolytic invasiveness of tumor cells and normal placental trophoblast cells. TIMP-3 may be involved in regulating trophoblastic invasion of the uterus as well as in regulating remodeling of the extracellular matrix during the folding of epithelia, and in the formation, branching and expansion of epithelial tubes.

### Supplied As

200ug/ml of Ab Purified from rabbit anti-serum by Protein A. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

AKT Signaling, Angiogenesis, Cardiovascular, Endothelial Cell Marker

## Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---