

TTF-1 / NKX2.1 (Thyroid & Lung Epithelial Marker) Antibody

Mouse Monoclonal Antibody [Clone 8G7G3/1]

Catalog No	Format	Size
7080-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
7080-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
7080-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	8G7G3/1
Gene Name	NKX2-1
Immunogen	Rat full length TTF-1 recombinant protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	40kDa
Cellular Localization	Nucleus
Species Reactivity	Human, Mouse, Rat
Positive Control	H441-4 or H345 cells. Normal thyroid or lung., MAD109, MLE-15

*Optimal dilution for a specific application should be determined.


Product Images for TTF-1 / NKX2.1 (Thyroid & Lung Epithelial Marker) Antibody



Formalin-fixed, paraffin-embedded human Thyroid stained with TTF-1 Mouse Monoclonal Antibody (8G7G3/1).



SDS-PAGE Analysis of Purified TTF-1 Mouse Monoclonal Antibody (8G7G3/1). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Lung Adenocarcinoma stained with TTF-1 Mouse Monoclonal Antibody (8G7G3/1).

Specificity & Comments

Recognizes a protein of 40kDa, identified as Thyroid transcription factor-1 (TTF-1). TTF-1 is a member of the NKx2 family of homeodomain transcription factors. It is expressed in epithelial cells of the thyroid gland and the lung. Nuclei from liver, stomach, pancreas, small intestine, colon, kidney, breast, skin, testes, pituitary, prostate, and adrenal glands are unreactive. Anti-TTF-1 is useful in differentiating primary adenocarcinoma of the lung from metastatic carcinomas originating in the breast, mediastinal germ cell tumors, and malignant mesothelioma. It can also be used to differentiate small cell lung carcinoma from lymphoid infiltrates. Loss of TTF-1 expression in non-small cell lung carcinoma has been associated with aggressive behavior of such neoplasms. TTF-1 reactivity is also seen in thyroid malignancies.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cancer, Neuroscience
