

SUMO-1 Antibody

Mouse Monoclonal Antibody [Clone SUMO1/1188]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 7341-MSM2-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 7341-MSM2-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 7341-MSM2-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|---------------------|---|
| Flow Cytometry (Flow) | 1-2ug/million cells | |
| Immunofluorescence (IF) | 1-3ug/ml | |
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details


| | |
|-------------------------------|--|
| Clone | SUMO1/1188 |
| Gene Name | SUMO1 |
| Immunogen | Recombinant human SUMO1 protein |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG1 / Kappa |
| Mol. Weight of Antigen | 11.5kDa (Monomer); 90kDa (Heteromer) |
| Cellular Localization | Cell membrane, Cytoplasm, Nucleus, Nucleus membrane, Nucleus speckle, PML body |
| Species Reactivity | Human, Rat |
| Positive Control | Breast carcinoma. |

*Optimal dilution for a specific application should be determined.


Product Images for SUMO-1 Antibody




Formalin-fixed, paraffin-embedded human Testicular Carcinoma stained with SUMO-1 Monoclonal Antibody (SUMO1/1188)




Formalin-fixed, paraffin-embedded human Tonsil stained with SUMO-1 Monoclonal Antibody (SUMO1/1188)



Formalin-fixed, paraffin-embedded human Tonsil stained with SUMO-1 MAb (SUMO1/1188)



Formalin-fixed, paraffin-embedded Rat Pancreas stained with SUMO-1 MAb (SUMO1/1188)



Formalin-fixed, paraffin-embedded human Ovarian Carcinoma stained with SUMO-1 Monoclonal Antibody (SUMO1/1188)

Specificity & Comments

This MAb is specific to SUMO-1 and shows no cross-reaction with either SUMO-2 or SUMO-3. The small ubiquitin-related modifier (SUMO) proteins, which include SUMO-1, SUMO-2 and SUMO-3, belong to the ubiquitin-like protein family. Like ubiquitin, the SUMO proteins are synthesized as precursor proteins that undergo processing before conjugation to target proteins. Also, both utilize the E1, E2, and E3 cascade enzymes for conjugation. However, SUMO and ubiquitin differ with respect to targeting. Ubiquitination predominantly targets proteins for degradation, whereas sumoylation targets proteins to a variety of cellular processing, including nuclear transport, transcriptional regulation, apoptosis and protein stability. The unconjugated SUMO-1 protein localizes to the nuclear membrane.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Autophagy, Cardiovascular, Cytokine Signaling, Immunology, Infectious Disease, Nuclear Marker, Transcription Factors