

CD106 / VCAM1 (Activated Endothelial Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone VCAM1/843]


Catalog No	Format	Size
7412-MSM2-P0	Purified Ab with BSA and Azide	200ug/ml
7412-MSM2-P1	Purified Ab with BSA and Azide	200ug/ml
7412-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Immunohistochemistry (IHC)	1-2ug/ml

Product Details	
Clone	VCAM1/843
Gene Name	VCAM1
Immunogen	Recombinant human VCAM1 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	110kDa
Cellular Localization	Membrane
Species Reactivity	Human
Positive Control	Human placenta or tonsil.

**Optimal dilution for a specific application should be determined.*

Product Images for CD106 / VCAM1 (Activated Endothelial Cell Marker) Antibody



Formalin-fixed, paraffin-embedded human Tonsil stained with CD106 Monoclonal Antibody (VCAM1/843).

Specificity & Comments

Recognizes a protein of 110kDa, identified as CD106 (also known as vascular cell adhesion molecule-1 (VCAM-1) and INCAM-100). CD106 is a member of the Ig superfamily of adhesion molecules and is expressed at high levels on cytokine stimulated vascular endothelial cells, and at minimal levels on un-stimulated endothelial cells. It is also present on follicular and inter-follicular dendritic cells of lymph nodes, myoblasts, and some macrophages. CD106 serves as a ligand for leukocyte integrin (VLA-4 or CD49d/CD29) and mediates cell adhesion of leukocytes to activated endothelium. It plays a role in various immunological and inflammatory responses.

Research Areas

BBB VCAM-1 Signaling, Cardiovascular, Cytokine Signaling, Endothelial Cell Marker, Hypoxia, Immunology, Mesenchymal Stem Cell Differentiation, Stem Cell Differentiation

Known Applications & Suggested Dilutions

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT)(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes)Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
