

Renal Cell Carcinoma / gp200 (Carbonic Anhydrase IX) Antibody

Mouse Monoclonal Antibody [Clone SPM314]

Catalog No	Format	Size
768-MSM1X-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
768-MSM1X-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
768-MSM1X-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	SPM314
Gene Name	CA9
Immunogen	Microsomal fraction of human renal cortical tissue homogenate
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	200kDa
Cellular Localization	Cell membrane, Cell projection, Microvillus membrane, Nucleolus, Nucleus
Species Reactivity	Horse, Human
Positive Control	Normal kidney or renal cell carcinoma.

**Optimal dilution for a specific application should be determined.*

Product Images for Renal Cell Carcinoma / gp200 (Carbonic Anhydrase IX) Antibody



SDS-PAGE Analysis of Purified Carbonic anhydrase 9 Mouse Monoclonal Antibody (SPM314). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human Renal Cell Carcinoma stained with RCC Monoclonal Antibody (SPM314).

Specificity & Comments

Recognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowman's capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This MAb may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Hypoxia, Angiogenesis
