

Recombinant Prostein (p501S) / Solute Carrier Family 45 Member 3 (SLC45A3) Antibody

Rabbit Monoclonal Antibody [Clone ZR9]

Catalog No	Format	Size
85414-RBM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
85414-RBM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
85414-RBM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	ZR9
Gene Name	SLC45A3
Immunogen	Synthesized peptides to the N terminus of human SLC45A3 (prostein) protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	59kDa
Cellular Localization	Membrane
Species Reactivity	Human
Positive Control	Normal prostate or prostate carcinoma.


*Optimal dilution for a specific application should be determined.

Product Images for Recombinant Prostein (p501S) / Solute Carrier Family 45 Member 3 (SLC45A3) Antibody



Formalin-fixed, paraffin-embedded human prostate carcinoma stained with Prostein (p501S) Recombinant Rabbit Monoclonal Antibody (ZR9). HIER: Tris/EDTA, pH9.0, 45min. 2 °: HRP-polymer, 30min. DAB, 5min.


Formalin-fixed, paraffin-embedded human prostate stained with Prostein (p501S) Recombinant Rabbit Monoclonal Antibody (ZR9). Inset: PBS instead of primary antibody; secondary only negative control.



SDS-PAGE Analysis Purified Prostein (p501S) Recombinant RabbitMonoclonal Antibody (ZR9). Confirmation of Purity and Integrity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteins using Prostein (p501S) Recombinant Rabbit Monoclonal Antibody (ZR9). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human prostate stained with Prostein (p501S) Recombinant Rabbit Monoclonal Antibody (ZR9). HIER: Tris/EDTA, pH9.0, 45min. 2 °: HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

Human prostein is a 553 aa protein identified by cDNA library subtraction and subsequent high throughput microarray screening of prostate cancer. Prostein has multiple transmembrane domains. Prostein has been shown to be uniquely expressed in normal and cancerous prostatic tissues. By immunohistochemistry, prostein is expressed in the vast majority of normal and malignant prostatic tissues, regardless of grade and metastatic status. No protein expression is detected in normal and malignant tissue samples representing the great majority of essential tissues and tumors. In particular, prostein is expressed in most of poorly differentiated prostatic carcinoma, including small cell prostate carcinoma. Prostein is more specific and sensitive for prostatic carcinomas than PSA and PSAP.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in CHO cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Mast Cell Marker