

CD1a (Mature Langerhans Cells Marker) Antibody

Mouse Monoclonal Antibody [Clone SPM120]

Catalog No	Format	Size
909-MSM1X-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
909-MSM1X-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
909-MSM1X-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	SPM120
Gene Name	CD1A
Immunogen	Human thymus cells
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	49kDa
Cellular Localization	Cell membrane, Endosome membrane, Membrane raft
Species Reactivity	Human
Positive Control	Jurkat, MOLT-4 cells. Paracortex in a tonsil or a reactive lymph node.

*Optimal dilution for a specific application should be determined.


Product Images for CD1a (Mature Langerhans Cells Marker) Antibody



Flow Cytometric Analysis of Jurkat cells using CD1a Monoclonal Antibody (SPM120) followed by goat anti-Mouse IgG-CF488 (Blue); Isotype control (Red).



SDS-PAGE Analysis of Purified T-cell surface glycoprotein CD1a Mouse Monoclonal Antibody (SPM120). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Skin stained with CD1a Monoclonal Antibody (SPM120).

Specificity & Comments

At least five CD1 genes (CD1a, b, c, d, and e) are identified. CD1 proteins have been demonstrated to restrict T cell response to non-peptide lipid and glycolipid antigens and play a role in non-classical antigen presentation. CD1a is a non-polymorphic MHC Class 1 related cell surface glycoprotein, expressed in association with Beta-2 microglobulin. Anti-CD1a labels Langerhans cell histiocytosis (Histiocytosis X), extranodal histiocytic sarcoma, a subset of T-lymphoblastic lymphoma/leukemia, and interdigitating dendritic cell sarcoma of the lymph node. When combined with antibodies against TTF-1 and CD5, anti-CD1a is useful in distinguishing between pulmonary and thymic neoplasms since CD1a is consistently expressed in thymic lymphocytes in both typical and atypical thymomas, but only focally in 1/6 of thymic carcinomas and not in lymphocytes in pulmonary neoplasms. Anti-CD1a is reported to be a new marker for perivascular epithelial cell tumor (PEComa).

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Dendritic Cell Marker, Immunology