

N myc (and STAT) Interactor / NMI Antibody

Mouse Monoclonal Antibody [Clone PCRP-NMI-1C1]



Catalog No	Format	Size
9111-MSM1-P0	Purified Ab with BSA and Azide	200ug/ml
9111-MSM1-P1	Purified Ab with BSA and Azide	200ug/ml
9111-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Flow Cytometry (Flow)	1-2ug/million cells

Product Details	
Clone	PCRP-NMI-1C1
Gene Name	NMI
Immunogen	Recombinant full-length human NMI protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b
Mol. Weight of Antigen	38kDa
Cellular Localization	Cytoplasm, Nucleus, Secreted
Species Reactivity	Human
Positive Control	HeLa or K562 cells.


*Optimal dilution for a specific application should be determined.

Product Images for N myc (and STAT) Interactor / NMI Antibody



SDS-PAGE Analysis of Purified NMI Mouse Monoclonal Antibody (PCRP-NMI-1C1). Confirmation of Purity and Integrity of Antibody.

Analysis of Protein Array containing more than 19,000 full-length human proteins using NMI Mouse Monoclonal Antibody (PCRP-NMI-1C1). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Flow Cytometric Analysis of PFA-fixed HeLa cells. NMI Mouse Monoclonal Antibody (PCRP-NMI-1C1) followed by goat anti-mouse IgG-CF488 (blue); unstained cells (red).

Specificity & Comments

Nmi (for N-Myc interactor) is an interferon inducible protein that associates with multiple transcription factors, including c-Myc, n-Myc, Max, and c-Fos, which contain bHLH-ZIP, bHLH, or Zip domains. Nmi is ubiquitously expressed at low levels throughout various fetal and adult tissues and at higher levels in myeloid leukemias and cell lines overexpressing c-Myc. In addition to binding Myc proteins, Nmi also associates with the Stat family of transcription factors, where it enhances Stat-dependent transcription. Although Nmi lacks an intrinsic DNA binding or activation domain, Nmi enhances the transcriptional activity of the Stat proteins, in response to cytokine stimulation, by recruiting the Stat1 and Stat5 transcriptional coactivators, CREB-binding protein (CBP) and p300. In vitro studies indicate that Nmi, expressed in conjunction with CBP, enhances the transcriptional responsiveness of Stat5 to IL-2 stimulation five-fold over CBP alone by increasing the affinity of Stat proteins for CBP/p300.

Research Areas

Cardiovascular, Infectious Disease, Nuclear Marker

Known Applications & Suggested Dilutions

Flow Cytometry (1-2ug/million cells) | ,Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.