

Recombinant CD22 / BL-CAM (B-Cell Marker) Antibody

Mouse Monoclonal Antibody [Clone rBLCAM/4108]

Catalog No	Format	Size
933-MSM8-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
933-MSM8-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
933-MSM8-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	rBLCAM/4108
Gene Name	CD22
Immunogen	Recombinant fragment (around aa52-178) of human BL-CAM/CD22 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	130-140kDa
Cellular Localization	Cell membrane
Species Reactivity	Human
Positive Control	lymph node, Ramos, Raji cells or human peripheral blood lymphocytes (hPBL). Human tonsil, Spleen.

**Optimal dilution for a specific application should be determined.*


Product Images for Recombinant CD22 / BL-CAM (B-Cell Marker) Antibody




IHC analysis of formalin-fixed, paraffin-embedded human brain. Negative tissue control using rBLCAM/4108 at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis of Purified CD22 Recombinant Mouse Monoclonal Antibody (rBLCAM/4108). Confirmation of Purity and Integrity of Antibody.



Western Blot analysis of Human Spleen and Ramos lysates using CD22 Recombinant Mouse Monoclonal Antibody. (rBLCAM/4108).



Formalin-fixed, paraffin-embedded human spleen stained with CD22 Recombinant Mouse Monoclonal Antibody (rBLCAM/4108).

Specificity & Comments

Recognizes a protein of 130-140kDa, identified as CD22 (also known as BL-CAM). CD22 expression is restricted to normal and neoplastic B cells and is absent from other haemopoietic cell types. In B-cell ontogeny, CD22 is first expressed in the cytoplasm of pro-B and pre-B cells, and on the surface as B cells mature to become IgD+. It is not expressed by plasma cells, CD22 is found highly expressed in follicular mantle and marginal zone B-cells, and while germinal center B-cells are relatively weak. CD22 is a member of the immunoglobulin superfamily and serves as an adhesion receptor for sialic acid-bearing ligands expressed on erythrocytes and all leukocyte classes. It also associates with tyrosine kinases and play a role in signal transduction and B-cell activation.

Supplied As

200ug/ml of Ab produced in CHO cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Immunology, B Cell Markers, Dendritic Cell Marker, Hematopoietic Stem Cells

Limitations and Warranty

This antibody is available for research use only and is not approved for diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.