

# CD163 (Monocyte & Macrophage Marker) Antibody

Mouse Monoclonal Antibody [Clone M130/2164]

Catalog No	Format	Size
9332-MSM4-P0	Purified Ab with BSA and Azide	200ug/ml
9332-MSM4-P1	Purified Ab with BSA and Azide	200ug/ml
9332-MSM4-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

# Applications

Immunohistochemistry (IHC)

1-2ug/ml

**Tested Dillution** 

#### **Product Details**

Clone	M130/2164
Gene Name	MVP
Immunogen	Recombinant fragment of human CD163 protein (around aa 43-196) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2b / Kappa
Mol. Weight of Antigen	140kDa
Cellular Localization	Cytoplasm, Nuclear pore complex, Nucleus, Perinuclear region
Species Reactivity	Human
Positive Control	HPBL cells. Skin, Lung, Placenta or Histiocytoma.

\*Optimal dilution for a specific application should be determined.

# Product Images for CD163 (Monocyte & Macrophage Marker) Antibody



Formalin-fixed, paraffin-embedded human Lymph Node in Colon stained with CD163 Mouse Monoclonal Antibody (M130/2164).



SDS-PAGE Analysis Purified CD163 Mouse Monoclonal Antibody (M130/2164). Confirmation of Purity and Integrity of Antibody.





Formalin-fixed, paraffin-embedded human Tonsil stained with CD163 Mouse Monoclonal Antibody (M130/2164).



Analysis of Protein Array containing more than 19,000 full-length human proteinsusing CD163 Mouse Monoclonal Antibody (M130/2164). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProtTM array. Z-scores are described in units of standard deviations (SD?s) above the mean value of all signals generated on that array. If targets on HuProtTM are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD?s) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human Lymph Node stained with CD163 Mouse Monoclonal Antibody (M130/2164).

#### **Specificity & Comments**

This MAb recognizes a protein of 140kDa, identified as CD163. It has been identified as an acute phase-regulated transmembrane protein whose function is to mediate the endocytosis of haptoglobinhemoglobin complexes. This receptor is expressed on the surface of monocytes with low expression and on tissue macrophages, histiocytes with high expression. Staining with anti-CD163 has been helpful to distinguish synovial macrophages from synovial intimal fibroblasts in rheumatoid arthritis, where its specificity for macrophages was found to be superior to that of anti-CD68. Increased levels of CD163 were also detected in patients with microbial infections and myelomonocytic leukemias. Anti-CD163 is of considerable value for selective identification of monocytes and macrophages at a certain stage of differentiation and is suitable for diagnosing myelomonocytic or monocytic leukaemia and neoplasms of true histiocytic origin. CD163 is positive in skin (histiocytes), gut, Kupffer cells, a few alveolar macrophages, macrophages in the placenta, and in macrophages in inflamed tissues including tumor tissue.

#### **Research Areas**

Cardiovascular, Dendritic Cell Marker, Hematopoietic Stem Cells, Infectious Disease

#### **Known Applications & Suggested Dilutions**

ELISA (For coating, order antibody without BSA) | ,Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues is enhanced by heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0 for 45 min at 95&degC followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.

#### **Supplied As**

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

# **Storage and Stability**

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.



### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

