

CD86 (Dendritic Cells Maturation Marker) Antibody

Mouse Monoclonal Antibody [Clone SPM600]

Catalog No	Format	Size
942-MSM2X-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
942-MSM2X-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
942-MSM2X-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunofluorescence (IF)	1-3ug/ml	

Product Details


Clone	SPM600
Gene Name	CD86
Immunogen	Recombinant human full-length CD86 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	70kDa
Cellular Localization	Cell membrane
Species Reactivity	Human
Positive Control	293T or Jukat cells. Monocytes and dendritic cells, activated T, B and natural killer cells in lymph node or tonsil.

*Optimal dilution for a specific application should be determined.

Product Images for CD86 (Dendritic Cells Maturation Marker) Antibody



SDS-PAGE Analysis of Purified T-lymphocyte activation antigen CD86 Mouse Monoclonal Antibody (SPM600). Confirmation of Purity and Integrity of Antibody.



Immunofluorescence staining of Ramos cells using CD86 Mouse Monoclonal Antibody (SPM600); followed by goat anti-Mouse IgG conjugated to CF488 (green). Nuclei are stained with Reddot.

Specificity & Comments

Recognizes a protein of 70kDa, which is identified as CD86. CD86 is a type I transmembrane glycoprotein and a member of the immunoglobulin superfamily of cell surface receptors. It is expressed at high levels on resting peripheral monocytes and dendritic cells and at very low density on resting B and T lymphocytes. CD86 expression is rapidly upregulated by B cell specific stimuli with peak expression at 18 to 42 hours after stimulation. CD86, along with CD80/B71, is an important accessory molecule in T cell co-stimulation via its interaction with CD28 and CD152/CTLA4. Since CD86 has rapid kinetics of induction, it is believed to be the major CD28 ligand expressed early in the immune response. It is also found on malignant Hodgkin and Reed Sternberg (HRS) cells in Hodgkin's disease.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

B Cell Markers, Cardiovascular, Cytokine Signaling, Dendritic Cell Marker, Hematopoietic Stem Cells, Immuno Oncology, Immunology, Infectious Disease, Signal Transduction

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.
