

CD44v6 (Marker of Tumor Metastasis) Antibody

Mouse Monoclonal Antibody [Clone 2F10]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 960-MSM11-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 960-MSM11-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 960-MSM11-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|-------------------------|---------------------|------|
| Flow Cytometry (Flow) | 1-2ug/million cells | |
| Immunofluorescence (IF) | 1-3ug/ml | |
| Western Blot (WB) | 2-4ug/ml | |

Product Details


| | |
|-------------------------------|---|
| Clone | 2F10 |
| Gene Name | CD44 |
| Immunogen | Recombinant fragment corresponding to the v9 domain of human CD44 |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG1 / Kappa |
| Mol. Weight of Antigen | 80-95kDa |
| Cellular Localization | Cell membrane, Cell projection, Microvillus |
| Species Reactivity | Human |
| Positive Control | Cervix or Tongue Squamous Cell Carcinoma |

*Optimal dilution for a specific application should be determined.


Product Images for CD44v6 (Marker of Tumor Metastasis) Antibody



Formalin-fixed, paraffin-embedded human Cervix Carcinoma stained with CD44v6 Mouse Monoclonal Antibody (2F10).



SDS-PAGE Analysis of Purified CD44v6 Mouse Monoclonal Antibody (2F10). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with CD44v6 Mouse Monoclonal Antibody (2F10).

Specificity & Comments

This antibody recognizes an epitope encoded by exon v6 on the variant portion of human CD44. The CD44 molecule belongs to a family of cellular adhesion molecules found on a wide range of normal and malignant cells in epithelial, mesothelial and hemopoiesis tissues. CD44 is a single gene with 20 exons, of which 10 are normally expressed to encode the basic CD44 (H-CAM) molecule. The additional 10 exons (v1 to v10) are only expressed by alternative splicing of the nuclear RNA. The expression of specific cell adhesion molecule CD44 splice variants has been reported to be associated with metastasis in certain human malignancies.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Articular Cartilage Extracellular Matrix, Cardiovascular, Cytokine Signaling, Hematopoietic Stem Cells, Immunology, Mesenchymal Stem Cell Differentiation, Neural Stem Cells