

## Recombinant CD44 / HCAM (Cancer Stem Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone HCAM/6459R]

Catalog No	Format	Size
960-RBM23-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
960-RBM23-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
960-RBM23-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	HCAM/6459R
<b>Gene Name</b>	CD44
<b>Immunogen</b>	Recombinant full-length human CD44 protein
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	80-95kDa
<b>Cellular Localization</b>	Cell membrane, Cell projection, Microvillus
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Paracortex in tonsil or lymph node (IHC). HeLa, A549

\*Optimal dilution for a specific application should be determined.


### Product Images for Recombinant CD44 / HCAM (Cancer Stem Cell Marker) Antibody




Formalin-fixed, paraffin-embedded human breast stained with CD44 Recombinant Rabbit Monoclonal Antibody (HCAM/6459R).




IHC analysis of formalin-fixed, paraffin-embedded human brain. Negative tissue control using HCAM/6459R at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.




SDS-PAGE Analysis of Purified CD44 Recombinant Rabbit Monoclonal Antibody (HCAM/6459R). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of HeLa cell lysate using CD44 Recombinant Rabbit Monoclonal Antibody (HCAM/6459R).



Western Blot Analysis of A549 and HeLa cell lysates using CD44 Recombinant Rabbit Monoclonal Antibody (HCAM/6459R).



Formalin-fixed, paraffin-embedded human esophagus stained with CD44 Recombinant Rabbit Monoclonal Antibody (HCAM/6459R).

### Specificity & Comments

Recognizes a cell surface glycoprotein of 80-95kDa (CD44) on lymphocytes, monocytes, and granulocytes (Leucocyte Typing Workshop V). Its epitope is resistant to digestion by trypsin and chymotrypsin. This MAb selectively interferes with lymphocyte binding to lymph node, mucosal and synovial endothelium. The CD44 family of glycoproteins exists in a number of variant isoforms, the most common being the standard 85-95kDa or hematopoietic variant (CD44s). Higher molecular weight isoforms are described in epithelial cells (CD44v), which are believed to function in intercellular adhesion and stromal binding. CD44 immunostaining is commonly used for the discrimination of urothelial transitional cell carcinoma in-situ from non-neoplastic changes in the urothelium.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Articular Cartilage Extracellular Matrix, Cardiovascular, Cytokine Signaling, Hematopoietic Stem Cells, Immunology, Mesenchymal Stem Cell Differentiation, Neural Stem Cells