

CD63 (Late Endosomes Marker) Antibody

Mouse Monoclonal Antibody [Clone LAMP3/2881]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 967-MSM11-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 967-MSM11-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 967-MSM11-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details


| | |
|-------------------------------|--|
| Clone | LAMP3/2881 |
| Gene Name | CD63 |
| Immunogen | Recombinant fragment (around aa 100-197) of human CD63 protein |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG2a / Kappa |
| Mol. Weight of Antigen | 26kDa (core protein); 30-60kDa (glycosylated) |
| Cellular Localization | Cell membrane, Cell surface, Endosome, Extracellular exosome, Late endosome membrane, Lysosome membrane, Melanosome, Multivesicular body, Secreted |
| Species Reactivity | Human |
| Positive Control | HL60, SK-MEL-28, THP-1 or NIH/3T3 cells. Spleen or Melanoma or lymphoma. |

*Optimal dilution for a specific application should be determined.



Product Images for CD63 (Late Endosomes Marker) Antibody



Western Blot Analysis of Jurkat cell lysate using CD63 Mouse Monoclonal Antibody (LAMP3/2881)




SDS-PAGE Analysis of Purified CD63 Mouse Monoclonal Antibody (LAMP3/2881). Confirmation of Purity and Integrity of Antibody.




Western Blot Analysis of human spleen tissue lysate using CD63 Mouse Monoclonal Antibody (LAMP3/2881)

Analysis of Protein Array containing >19,000 full-length human proteins using CD63 Mouse Monoclonal Antibody (LAMP3/2881) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Western blot analysis of human spleen, mouse spleen, rat spleen, hamster spleen and guinea pig spleen tissue lysates using CD63 Mouse Monoclonal Antibody (LAMP3/2881).



Formalin-fixed, paraffin-embedded human Melanoma stained with CD63 Mouse Monoclonal Antibody (LAMP3/2881)

Specificity & Comments

This MAb recognizes protein of 26kDa-60kDa, which is identified as CD63. The tetraspanins are integral membrane proteins expressed on cell surface and granular membranes of hematopoietic cells and are components of multi-molecular complexes with specific integrins. The tetraspanin CD63 is a lysosomal membrane glycoprotein that translocates to the plasma membrane after platelet activation. CD63 is expressed on activated platelets, monocytes and macrophages, and is weakly expressed on granulocytes, T cell and B cells. It is located on the basophilic granule membranes and on the plasma membranes of lymphocytes and granulocytes. CD63 is a member of the TM4 superfamily of leukocyte glycoproteins that includes CD9, CD37 and CD53, which contain four transmembrane regions. CD63 may play a role in phagocytic and intracellular lysosome-phagosome fusion events. CD63 deficiency is associated with Hermansky-Pudlak syndrome and is strongly expressed during the early stages of melanoma progression.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Immunology