

CD68 (Macrophage Marker) Antibody

Mouse Monoclonal Antibody [Clone KP1 + C68/684]

Catalog No	Format	Size
968-MSM3-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
968-MSM3-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
968-MSM3-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	KP1 + C68/684
Gene Name	CD68
Immunogen	Subcellular fraction of human alveolar macrophages (KP1); Recombinant human CD68 protein (C68/684)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	110kDa
Cellular Localization	Cell membrane, Endosome membrane, Lysosome membrane
Species Reactivity	Cat, Human, Monkey, Mouse, Rat
Positive Control	lymph node or spleen., Tonsil

*Optimal dilution for a specific application should be determined.


Product Images for CD68 (Macrophage Marker) Antibody



Formalin-fixed, paraffin-embedded human Tonsil stained with CD68 Mouse Monoclonal Antibody (KP1+C68/684).



SDS-PAGE Analysis of Purified Macrosialin Mouse Monoclonal Antibody (KP1+C68/684). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Histiocytoma stained with CD68 Mouse Monoclonal Antibody (KP1+C68/684).

Specificity & Comments

This antibody recognizes a glycoprotein of 110kDa, which is identified as CD68. It is important for identifying macrophages in tissue sections. It stains macrophages in a wide variety of human tissues, including Kupffer cells and macrophages in the red pulp of the spleen, in lamina propria of the gut, in lung alveoli, and in bone marrow. It reacts with myeloid precursors and peripheral blood granulocytes. It also reacts with plasmacytoid T cells, which are supposed to be of monocyte/macrophage origin. It shows strong granular cytoplasmic staining of chronic and acute myeloid leukemia and also reacts with rare cases of true histiocytic neoplasia. Lymphomas are negative or show few granules.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Hematopoietic Stem Cells, Immunology
