

Recombinant CD81 / TAPA-1 Antibody

Rabbit Monoclonal Antibody [Clone C81/2885R]

Catalog No	Format	Size
975-RBM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
975-RBM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
975-RBM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	C81/2885R
Gene Name	CD81
Immunogen	B-Cell line derived from a Burkitt lymphoma
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	26kDa
Cellular Localization	Basolateral cell membrane, Cell membrane
Species Reactivity	Human, Mouse, Rat
Positive Control	A293, A431, and human lymphocytes. Tonsil or Liver., Daudi, HL-60, HUT-102, HUT-78, Jurkat, Kg1a, MCF-7, MG63, Raji, Ramos, U266, Human Brain

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant CD81 / TAPA-1 Antibody




Flow cytometry analysis of MCF-7 cells unstained (gray) or stained with CD81 Rabbit Recombinant Monoclonal Antibody (C81/2885R) and goat anti-mouse CF647 (red).




Flow cytometry analysis of bead-bound exosomes derived from MCF-7 cells. Unstained exosomes (gray) or exosomes stained with CF568-labeled CD81 monoclonal antibody (2885R) (orange).



Western Blot Analysis of Human Brain tissue lysate using CD81 Recombinant Rabbit Monoclonal Antibody (C81/2885R).



SDS-PAGE Analysis of Purified CD81 antigen Recombinant Rabbit Monoclonal Antibody (C81/2885R). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human Prostate Carcinoma stained with CD81 Rabbit Recombinant Monoclonal Antibody (C81/2885R).

Specificity & Comments

Recognizes a protein of 26kDa, identified as CD81 (Workshop VI; Code CD81.1)). CD81 has a very broad cellular distribution, being expressed on T- and B-lymphocytes, NK cells, thymocytes, eosinophils, fibroblasts, epithelial and endothelial cells. Neutrophils, erythrocytes and platelets are negative, while monocytes are variably positive. CD81 is a member of a family of tetraspan transmembrane proteins, including CD9, CD37, CD53, CD63, and CD82. It associates with CD19, CD21, Leu 13, and integrins on cell membrane and is involved in signal transduction in B lymphocyte development and cell adhesion. CD81 also acts as a receptor for the envelope protein E2 of chronic hepatitis C virus. Antibodies to CD81 have anti-proliferative effects on different lymphoid cell lines, particularly those derived from large cell lymphomas.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Immunology, Complement System

Limitations and Warranty

This antibody is available for research use only and is not approved for diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.