

## Recombinant TNFS15 / VEGI (Vascular Endothelial Growth Inhibitor) Antibody

Rabbit Monoclonal Antibody [Clone VEGI/2052R]

Catalog No	Format	Size
9966-RBM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
9966-RBM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
9966-RBM2-P1ABX	Purified Ab WITHOUT BSA at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details


<b>Clone</b>	VEGI/2052R
<b>Gene Name</b>	TNFSF15
<b>Immunogen</b>	Recombinant full-length human VEGI protein
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	22kDa
<b>Cellular Localization</b>	Membrane, Secreted
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Endothelial cells. Colon, intestine, Kidney, Liver, Lung, Pancreas, Placenta, spleen and prostate.

*\*Optimal dilution for a specific application should be determined.*



### Product Images for Recombinant TNFS15 / VEGI (Vascular Endothelial Growth Inhibitor) Antibody



Formalin-fixed, paraffin-embedded human Parathyroid Mass stained with VEGI Rabbit Recombinant Monoclonal Antibody (VEGI /2052R).




Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with VEGI Rabbit Recombinant Monoclonal Antibody (VEGI /2052R).



SDS-PAGE Analysis of Purified Tumor necrosis factor ligand superfamily member 15 Recombinant Rabbit Monoclonal Antibody (VEGI/2052R). Confirmation of Purity and Integrity of Antibody.

Analysis of Protein Array containing more than 19,000 full-length human proteins using VEGI Rabbit Recombinant Monoclonal Antibody (VEGI/2052R). Z- and S-Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human Spleen stained with VEGI Rabbit Recombinant Monoclonal Antibody (VEGI /2052R).

### Specificity & Comments

VEGI is an anti-angiogenic cytokine that belongs to tumor necrosis factor superfamily, member 15 (TNFSF15). This protein is abundantly expressed in endothelial cells, but is not expressed in either B or T cells. The expression of this protein is inducible by TNF and IL-1 alpha. This cytokine is a ligand for receptor TNFRSF25 and decoy receptor TNFRSF21/DR6. It can activate NF-kappaB and MAP kinases, and acts as an autocrine factor to induce apoptosis in endothelial cells. This cytokine is also found to inhibit endothelial cell proliferation, and thus may function as an angiogenesis inhibitor. Reduced expression of VEGI has been reported as a marker of poor prognosis in breast cancer.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab purified by Protein A Column. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cytokine Signaling, Immunology