

E-Cadherin (CDH1) / CD324 (Intercellular Junction Marker) Antibody

Mouse Monoclonal Antibody [Clone CDH1/3256]

| Catalog No | Format | Size |
|-----------------|---|--------|
| 999-MSM14-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 999-MSM14-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 999-MSM14-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|---------------------|---|
| Flow Cytometry (Flow) | 1-2ug/million cells | |
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |

Product Details


| | |
|-------------------------------|---|
| Clone | CDH1/3256 |
| Gene Name | CDH1 |
| Immunogen | Recombinant fragment of human CDH1 protein (around aa 567-691) (exact sequence is proprietary) |
| Host | Mouse |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG1 / Kappa |
| Mol. Weight of Antigen | 120-80kDa (Mature); 135kDa (Precursor) |
| Cellular Localization | Adherens junction, Cell junction, Cell membrane, Endosome, Golgi apparatus, trans-Golgi network |
| Species Reactivity | Human |
| Positive Control | HT29 or SK-BR3 cells. Prostate or Colon carcinomas., LS174T, MCF-7, Raji |

*Optimal dilution for a specific application should be determined.



Product Images for E-Cadherin (CDH1) / CD324 (Intercellular Junction Marker) Antibody



Flow Cytometric Analysis of human trypsinized MCF-7 cells using E-Cadherin Mouse Monoclonal Antibody (CDH1/3256) followed by Goat anti-Mouse IgG-CF488 (Blue); Isotype control (Red)



SDS-PAGE Analysis of Purified E-Cadherin Mouse Monoclonal Antibody (CDH1/3256). Confirmation of Integrity and Purity of Antibody.



Formalin-fixed, paraffin-embedded human Breast Carcinoma stained with E-Cadherin Mouse Monoclonal Antibody (CDH1/3256).

Analysis of Protein Array containing more than 19,000 full-length human proteins using E-Cadherin Mouse Monoclonal Antibody (CDH1/3256) Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD, σ) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD, σ) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Specificity & Comments

Recognizes a protein of 120-80kDa, identified as E-cadherin. Cadherins comprise a family of Ca^{2+} -dependent adhesion molecules that function to mediate cell-cell binding critical to the maintenance of tissue structure and morphogenesis. The classical cadherins, E-, N- and P-cadherin, consist of large extracellular domains characterized by a series of five homologous NH₂ terminal repeats. The relatively short intracellular domains interact with a variety of cytoplasmic proteins, such as β -catenin, to regulate cadherin function. E-cadherin plays an important role in epithelial cell adhesion. A decreased expression of E-cadherin is associated with metastatic potential and poor prognosis in breast cancer, prostate and esophageal cancer. In combination with p120 Catenin, it is useful for the differentiation between ductal (E-cadherin +) and lobular (E-cadherin -) breast carcinomas. It may also help in diagnosis of mesothelioma.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Bladder Cancer, Cardiovascular, Colon Cancer, Developmental Biology, Immunology, Infectious Disease, Signal Transduction