



## Cytokeratin, Multi Antibody

Mouse Monoclonal Antibody [Clone SPM583]

Catalog No	Format	Size
MSM1X-393-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
MSM1X-393-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
MSM1X-393-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	SPM583
<b>Immunogen</b>	Keratin-enriched preparation from cultured human A431
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	Multiple
<b>Species Reactivity</b>	Cow, Frog, Goat, Guinea Pig, Human, Marmoset, Mouse, Pig, Rat
<b>Positive Control</b>	A431 cells, Colon carcinoma., Skin

\*Optimal dilution for a specific application should be determined.

### Product Images for Cytokeratin, Multi Antibody



SDS-PAGE Analysis of Purified Keratin, type II cytoskeletal 6A Mouse Monoclonal Antibody (SPM583). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded colon (10X) stained with Multi Keratin Monoclonal Antibody (SPM583).

### Specificity & Comments

This MAbs recognizes cytokeratin 4, 5, 6, 8, 10, 13, and 18. This is a broad-spectrum antibody which has been reported to differentiate epithelial tumors from non-epithelial tumors. Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis.

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

## Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---