

## Phosphotyrosine (P-Tyr) Antibody

Mouse Monoclonal Antibody [Clone PY265]

| Catalog No     | Format                                        | Size   |
|----------------|-----------------------------------------------|--------|
| MSM3-265-P0    | Purified Ab with BSA and Azide at 200ug/ml    | 20 ug  |
| MSM3-265-P1    | Purified Ab with BSA and Azide at 200ug/ml    | 100 ug |
| MSM3-265-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications            | Tested Dillution    | Note |
|-------------------------|---------------------|------|
| Flow Cytometry (Flow)   | 1-2ug/million cells |      |
| Immunofluorescence (IF) | 1-3ug/ml            |      |
| Western Blot (WB)       | 2-4ug/ml            |      |

### Product Details

|                               |                                                |
|-------------------------------|------------------------------------------------|
| <b>Clone</b>                  | PY265                                          |
| <b>Immunogen</b>              | Phosphotyrosine conjugated to BSA              |
| <b>Host</b>                   | Mouse                                          |
| <b>Clonality</b>              | Monoclonal                                     |
| <b>Isotype / Light Chain</b>  | IgG1 / Kappa                                   |
| <b>Mol. Weight of Antigen</b> | Depends upon the phosphorylated target         |
| <b>Cellular Localization</b>  | N/A                                            |
| <b>Species Reactivity</b>     | All species                                    |
| <b>Positive Control</b>       | MCF-7, MDA-231, T47-D cells. Breast carcinoma. |

\*Optimal dilution for a specific application should be determined.

### Product Images for Phosphotyrosine (P-Tyr) Antibody



SDS-PAGE Analysis of Purified Phosphotyrosine Mouse Monoclonal Antibody (PY265). Confirmation of Purity and Integrity of Antibody.

### Specificity & Comments

Protein phosphorylation is a fundamental event in the regulation of a large number of intracellular processes. Phosphorylation of specific tyrosine residues is the result of activation or stimulation of their respective protein tyrosine kinases. The phosphorylated proteins can be auto-phosphorylated kinases or certain cellular protein substrates. Tyrosine-phosphorylated proteins are involved in signal transduction and in the regulation of cell proliferation. Antibody to phosphotyrosine provides an excellent tool for the detection, characterization, and purification of phosphotyrosine containing proteins. This MAb shows no cross-reaction with other phosphoamino acids and is superb for multiple applications including staining of formalin/paraffin tissues.

### Supplied As

200ug/ml of Ab Purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.