

Nuclear Antigen (Pan-NuclearMarker) Antibody

Mouse Monoclonal Antibody [Clone NM106]


Catalog No	Format	Size
MSM4-106-P0	Purified Ab with BSA and Azide	200ug/ml
MSM4-106-P1	Purified Ab with BSA and Azide	200ug/ml
MSM4-106-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml

Applications	Tested Dillution
Flow Cytometry (Flow)	1-2ug/million cells
Immunofluorescence (IF)	1-3ug/ml
Immunohistochemistry (IHC)	1-2ug/ml

Product Details	
Clone	NM106
Gene Name	N/A
Immunogen	Nuclei of HL60 cells
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	Not Known
Cellular Localization	N/A
Species Reactivity	Human, Mouse, Rat
Positive Control	All human cells. Tonsil

**Optimal dilution for a specific application should be determined.*


Product Images for Nuclear Antigen (Pan-NuclearMarker) Antibody




Formalin-fixed, paraffin-embedded human tonsil stained with Pan-Nuclear Antigen Monoclonal Antibody (NM106).




Formalin-fixed, paraffin-embedded rat colon stained with Pan-Nuclear Antigen Monoclonal Antibody (NM106).




Formalin-fixed, paraffin-embedded tissue stained with Pan-Nuclear Antigen Monoclonal Antibody (NM106).



Immunofluorescent staining of PFA-fixed HeLa cells with Pan-Nuclear Antigen Monoclonal Antibody (NM106) followed by goat anti-mouse IgG-CF488 (green). Membranes labeled with phalloidin (red).



SDS-PAGE Analysis Purified Pan-Nuclear Antigen Monoclonal Antibody (NM106). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human tonsil stained with Pan-Nuclear Antigen Monoclonal Antibody (NM106).

Specificity & Comments

This MAb is an excellent marker for all nuclei in cells in tissues. It is a part of a new panel of reagents, which recognizes subcellular organelles or compartments of cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This MAb recognizes an antigen associated with the nuclei in all cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in subcellular fractions. It produces a speckled pattern in normal and malignant cells and may be used to stain the nuclei of cells in fixed or frozen tissue sections. It can also be used with paraformaldehyde fixed frozen tissue or cell preparations.

Known Applications & Suggested Dilutions

Flow Cytometry (1-2ug/million cells) | Immunofluorescence (1-2ug/ml) | Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT), (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95 & degC followed by cooling at RT for 20 minutes), Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.