

## Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) Antibody

Mouse Monoclonal Antibody [Clone SPM116]

Catalog No	Format	Size
MSM5X-257-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
MSM5X-257-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
MSM5X-257-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details

<b>Clone</b>	SPM116
<b>Gene Name</b>	KRT76
<b>Immunogen</b>	Human epidermal keratin
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 52kDa (CK8)
<b>Species Reactivity</b>	Chicken, Cow, Dog, Human, Monkey, Mouse, Rabbit, Rat
<b>Positive Control</b>	Adenocarcinomas, Skin

\*Optimal dilution for a specific application should be determined.

### Product Images for Cytokeratin, Basic (Type II or HMW) (Epithelial Marker) Antibody



SDS-PAGE Analysis of Purified Keratin, type II cytoskeletal 2 oral Mouse Monoclonal Antibody (SPM116). Confirmation of Purity and Integrity of Antibody.

Formalin-paraformaldehyde human Breast Carcinoma stained with Cytokeratin, HMWMAb (SPM116).

### **Specificity & Comments**

This MAb recognizes basic (Type II or HMW) cytokeratins, which include 67kDa (CK1); 64kDa (CK3); 59kDa (CK4); 58kDa (CK5); 56kDa (CK6); 52kDa (CK8). Twenty human keratins are resolved with two-dimensional gel electrophoresis into acidic (pI 6.0) subfamilies. The acidic keratins have molecular weights (MW) of 56.5, 55, 51, 50, 50', 48, 46, 45, and 40kDa. This MAb recognizes the 65-67, 64, 59, 58, 56, and 52kDa keratins of basic subfamily. Many studies have shown the usefulness of keratins as markers in cancer research and tumor diagnosis. SPM115/SPM116 is a broad spectrum anti pan-keratin antibody cocktail, which differentiates epithelial tumors from non-epithelial tumors e.g. squamous vs. adenocarcinoma of the lung, liver carcinoma, breast cancer, and esophageal cancer.

---

### **Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

### **Supplied As**

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

### **Storage and Stability**

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---

### **Research Areas**

Autophagy, Developmental Biology

---