

Recombinant Nuclear Antibody

Mouse Monoclonal Antibody [Clone rNM106]



Catalog No	Format	Size
MSM6-3466-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
MSM6-3466-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
MSM6-3466-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details	
Clone	rNM106
Immunogen	isolated nuclei of HL-60 cells
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Species Reactivity	Human
Positive Control	All human cells. Tonsil. Heart, Intestine, Liver Skeletal muscle.



*Optimal dilution for a specific application should be determined.

Product Images for Recombinant Nuclear Antibody



SDS-PAGE Analysis of Purified Pan-Nuclear Antigen Recombinant Mouse Monoclonal (rNM106). Confirmation of Purity and Integrity of Antibody.

Western Blot Analysis of Human Heart, Human Intestine, Human Liver and Human Skeletal muscle tissue lysates using Nuclear Antigen Recombinant Mouse Monoclonal Antibody (rNM106).



Western Blot Analysis of Heart, Liver and Skeletal muscle tissue lysates of different species using Nuclear Antigen Recombinant Mouse Monoclonal Antibody (rNM106).

Formalin-fixed, paraffin-embedded human tonsil stained with Pan-Nuclear Antigen Recombinant Mouse Monoclonal Antibody (rNM106).

Specificity & Comments

This MAb is an excellent marker for all nuclei in cells in tissues. It is a part of a new panel of reagents, which recognizes subcellular organelles or compartments of cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This MAb recognizes an antigen associated with the nuclei in all cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in subcellular fractions. It produces a speckled pattern in normal and malignant cells and may be used to stain the nuclei of cells in fixed or frozen tissue sections. It can also be used with paraformaldehyde fixed frozen tissue or cell preparations.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.