

## p27Kip1 (Mitotic Inhibitor/Suppressor Protein) Antibody

Mouse Monoclonal Antibody [Clone SX53G8]

Catalog No	Format	Size
1027-MSM1-CF488-100T	Purified Ab conjugated to CF488	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	
Immunofluorescence (IF)	1-3ug/ml	

### Product Details


<b>Clone</b>	SX53G8
<b>Gene Name</b>	CDKN1B
<b>Immunogen</b>	Purified GST-p27 fusion protein of human origin
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	25-26kDa
<b>Cellular Localization</b>	Nucleus
<b>Species Reactivity</b>	Human, Monkey, Mouse, Rat
<b>Positive Control</b>	Tonsil, breast or colon carcinoma. MCF7, MDA-MB-231, SK-BR-3 or T47D cells.

\*Optimal dilution for a specific application should be determined.


### Product Images for p27Kip1 (Mitotic Inhibitor/Suppressor Protein) Antibody



SDS-PAGE Analysis of Purified p27 Mouse Monoclonal Antibody (SX53G8). Confirmation of purity and integrity.



Immunofluorescence Analysis of PFA-fixed MCF-7 cells labeling p27 with p27 Mouse Monoclonal Antibody (SX53G8) followed by Goat anti-Mouse IgG-CF488 (Green). Membrane is stained with Phalloidin-CF640



Flow Cytometry of human p27 on HeLa cells. Black: cells alone; Grey: Isotype Control; Green: CF488-labeled p27 Monoclonal Antibody (SX53G8).

### Specificity & Comments

This MAb recognizes a 27kDa protein, identified as the p27Kip1, a cell cycle regulatory mitotic inhibitor. It is highly specific and shows no cross-reaction with other related mitotic inhibitors. In Western blotting of cell lysates from 7 human breast cancer cell lines (ZR75-1, ZR75-30, MCF-7, MDAMB453, T47D, CAL51, 734B), the antibody labels a single band corresponding to p27Kip1. It functions as a negative regulator of G1 progression and has been proposed to function as a possible mediator of TGF- induced G1 arrest. p27Kip1 is a candidate tumor suppressor gene. Reportedly, low p27 expression has been associated with unfavorable prognosis in renal cell carcinoma, colon carcinoma, breast carcinomas, non-small-cell lung carcinoma, hepatocellular carcinoma, multiple myeloma, and lymph node metastases in papillary carcinoma of the thyroid, as well as a more aggressive phenotype in carcinoma of the cervix.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

Antibody purified from Bioreactor Concentrate by Protein A/G and conjugated to various reporter molecules. Prepared in 10mM PBS with 0.05% BSA and 0.05% azide. Contact us if you require this Ab in a different format.

### Storage and Stability

Antibody with azide - store at 4 to 8°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cancer, Cardiovascular, Hypoxia, Immunology, AKT Signaling, Cytokine Signaling, Infectious Disease, Lung Cancer, Nuclear Marker, Ovarian Cancer, Signal Transduction, Transcription Factors