

## Recombinant TSH-Receptor, A-Chain (Thyroid Marker) Antibody

Rabbit Monoclonal Antibody [Clone TSHRB/9215R]

Catalog No	Format	Size
7253-RBM12-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
7253-RBM12-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
7253-RBM12-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	TSHRB/9215R
<b>Immunogen</b>	Recombinant full-length human TSH Receptor, B-Chain protein
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	86.84kDa
<b>Cellular Localization</b>	Basolateral cell membrane, Cell membrane
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	293T cells. Human thyroid gland. Heart, Skeletal muscle, kidney

\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant TSH-Receptor, A-Chain (Thyroid Marker) Antibody



Western blot analysis of Human Heart, Human Kidney and Human Skeletal muscle tissue lysates using TSHRB Recombinant Rabbit Monoclonal Antibody (TSHRB/9215R).



SDS-PAGE Analysis Purified TSHRB Recombinant Rabbit Monoclonal Antibody (TSHRB/9215R). Confirmation of Purity and Integrity of Antibody.

### Specificity & Comments

Thyroid-stimulating hormone (TSH, also known as thyrotropin) is a glycoprotein involved in the control of thyroid structure and metabolism, which stimulates the release of the thyroid hormones. TSH is regulated by thyroid hormone (T3) and various retinoid compounds. TSH binds to the thyroid-stimulating hormone receptor (TSHR), which is cleaved into two subunits, A and B, and plays a major role in regulating thyroid function. The third cytoplasmic loop of TSHR has been identified as critical for its role in regulating inositol phosphate and cAMP formation. In Grave s disease, an autoimmune disorder, TSHR is activated by autoantibodies, which may be stimulated by the cleavage of the A and B subunits.

### Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

## Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---