

Recombinant RCAS1 / Estrogen Receptor Binding Site Associated, Antigen 9 Antibody

Mouse Monoclonal Antibody [Clone rEBAG9/7264]

Catalog No	Format	Size
9166-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
9166-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
9166-MSM4-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	rEBAG9/7264
Immunogen	Recombinant full-length human RCAS1 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	24.38kDa
Cellular Localization	Golgi apparatus membrane
Species Reactivity	Human
Positive Control	Jurkat or HEK-293 cell lysates. Human skeletal muscle tissue. MCF-7, Ramos

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant RCAS1 / Estrogen Receptor Binding Site Associated, Antigen 9 Antibody



SDS-PAGE Analysis Purified RCAS1 Recombinant Mouse Monoclonal Antibody (rEBAG9/7264). Confirmation of Purity and Integrity of Antibody.



Formalin-fixed, paraffin-embedded human brain stained with RCAS1 Recombinant Mouse Monoclonal Antibody (rEBAG9/7264). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

EBAG9, also known as RCAS1, is an estrogen-transcribed protein. Soluble and membranous RCAS1 proteins may play a role in the immune escape of tumor cells by promoting T lymphocyte inhibition of growth and apoptosis. RCAS1 is expressed in a wide variety of cancers, including uterine, ovarian, and lung cancer cells, and acts as a ligand for a putative receptor present on peripheral lymphocytes. RCAS1 is highly expressed not only in cancer cells but also in non-tumor bile duct cells subject to immune attack. RCAS1 inhibits the in vitro growth of receptor-expressing cells and induces apoptosis, contributing to the ability of tumor cells to evade host immune surveillance. High expression of RCAS1 significantly correlates with tumor progression and with poor outcome for many cancer patients.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
