Recombinant CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody

Rabbit Monoclonal Antibody [Clone LFA2/8681R]

Catalog No	Format	Size
914-RBM22-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
914-RBM22-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
914-RBM22-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	LFA2/8681R
Immunogen	Recombinant fragment (around aa272-374) of the human CD2 protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	39.45kDa
Cellular Localization	Cell membrane
Species Reactivity	Human
Positive Control	Humde.an tonsil|spleen or lymph node, Jurkat

**Optimal dilution for a specific application should be determined.*


Product Images for Recombinant CD2 / Lymphocyte Function Antigen 2 (LFA-2) Antibody



Formalin-fixed, paraffin-embedded human spleen stained with CD2 Recombinant Rabbit Monoclonal Antibody (LFA2/8681R). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



Western blot analysis of Jurkat cell lysate using CD2 Recombinant Rabbit Monoclonal Antibody (LFA2/8681R).



SDS-PAGE Analysis of Purified CD2 Recombinant Rabbit Monoclonal Antibody (LFA2/8681R). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

CD2 is mainly present on the surface of mature T cells and NK cell membranes, while B cells are usually not expressed. It can be used to label the diagnosis and study of normal T cells and their associated tumors, such as pre-t cell lymphoma, marginal T cell lymphoma and anaplastic cell lymphoma. CD2 antibody is useful for identification of precursor and mature T-cell lymphomas. Aberrant loss of CD2 in T-cell lymphomas may help to distinguish them from reactive T-cell proliferations.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.