

## ICOS-L / ICOS Ligand / B7RP-1 (Immuno-Oncology Target) Antibody

Mouse Monoclonal Antibody [Clone ICOSL/7700]

Catalog No	Format	Size
23308-MSM9-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
23308-MSM9-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
23308-MSM9-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	ICOSL/7700
<b>Gene Name</b>	ICOSLG
<b>Immunogen</b>	Recombinant fragment of human ICOSLG (B7RP-1) protein (around aa 1-300) (exact sequence is proprietary)
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2b / Kappa
<b>Mol. Weight of Antigen</b>	36kDa
<b>Cellular Localization</b>	Cell Surface. Cytoplasm.
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	HL60 HuT78 or HUVEC lysates (WB). Human tonsil tissue (IHC). U937. Human Kidney, Human Spleen, K-562, Raji or Ramos

\*Optimal dilution for a specific application should be determined.


### Product Images for ICOS-L / ICOS Ligand / B7RP-1 (Immuno-Oncology Target) Antibody




Formalin-fixed, paraffin-embedded human colon carcinoma stained with ICOSL Mouse Monoclonal Antibody (ICOSL/7700). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis of Purified ICOS ligand Mouse Monoclonal Antibody (ICOSL/7700). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of Human Kidney, Human Spleen, K-562, Raji and Ramos lysates using ICOS-L Mouse Monoclonal Antibody (ICOSL/7700).



Formalin-fixed, paraffin-embedded human spleen stained with ICOS-L Mouse Monoclonal Antibody (ICOSL/7700). Inset: PBS instead of primary antibody; secondary only negative control.

### Specificity & Comments

This MAb recognizes a protein of 36kDa, identified as ICOS-L. It is a ligand for the ICOS receptor that initiates T and B cell proliferation and cytokine secretion. ICOS-L interactions play an essential role in T cell-dependent B cell activation in peripheral lymphoid organs such as spleen and lymph nodes. ICOS-L protein is present in myeloid leukocytes, and by Northern blot there are 2.4, 3.0, and 7.0 kb transcripts in brain, heart, kidney, and liver, with lower expression in colon and thymus, and a 1.1 kb transcript in leukocytes. Tumor necrosis factor alpha (TNF $\alpha$ ), granulocyte-macrophage colony-stimulating factor (GM-CSF) and interleukin-4 (IL-4) enhance its expression. LPS-induced up-regulation of ICOS-L is dependent on the MyD88-dependent signaling pathway.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cardiovascular, Immunology