Catalog # 10630-RBM8
Product Datasheet

Recombinant Podoplanin (PDPN) (Lymphatic Endothelial & Mesothelial Marker) Antibody

Rabbit Monoclonal Antibody [Clone PDPN/7808R]

Catalog No	Format	Size
10630-RBM8-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
10630-RBM8-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
10630-RBM8-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	PDPN/7808R
Gene Name	PDPN
Immunogen	Recombinant fragment (around aa1-100) of human PDPN protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	17kDa
Cellular Localization	Cell surface.
Species Reactivity	Human
Positive Control	cervical or lung carcinoma tissues.|Human tonsil

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant Podoplanin (PDPN) (Lymphatic Endothelial & Mesothelial Marker) Antibody



Western blot analysis of human Lung tissue lysate using PDPN Recombinant Rabbit Monoclonal Antibody (PDPN/7808R).



Formalin-fixed, paraffin-embedded human tonsil stained with Podoplanin Recombinant Rabbit Monoclonal Antibody (PDPN/7808R). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

It recognizes a muco-protein of 38-43kDa, which is identified as Podoplanin (PDPN). It localizes in stromal cells of peripheral lymphoid tissue and thymic epithelial cells. As a regulator of the lymphatic endothelium, podoplanin probably plays a role in maintaining the unique shape of podocytes. It is selectively expressed in lymphatic endotheliumÂ ?as well as lymphoangiomas,Â ?Kaposi sarcomas,Â ?and in a subset of angiosarcomas with probable lymphatic differentiation. Recent studies have also shown podoplanin to be a highly sensitive and relatively specific marker for epithelioid mesothelioma. Therefore, it can be used in a panel to distinguish mesotheliomas or mesothelial cells from pulmonary carcinomas.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Angiogenesis, Dendritic Cell Marker, Endothelial Cell Marker

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.