

Recombinant OLIG2 (Marker of Glial Brain Tumors) Antibody

Rabbit Monoclonal Antibody [Clone OLIG2/7366R]

Catalog No	Format	Size
10215-RBM6-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
10215-RBM6-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
10215-RBM6-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	OLIG2/7366R
Gene Name	OLIG2
Immunogen	Recombinant fragment (around aa200-300) of human OLIG2 protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	32kDa
Cellular Localization	Nucleus. Cytoplasm.
Species Reactivity	Human
Positive Control	THP-1 cells. Human brain or astrocytoma.

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant OLIG2 (Marker of Glial Brain Tumors) Antibody



IHC analysis of FFPE human tonsil (RNA expression: 0.0 nTPM). Negative tissue control using OLIG2 Recombinant Rabbit Monoclonal (OLIG2/7366R). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



SDS-PAGE Analysis of Purified Oligodendrocyte transcription factor 2 Recombinant Rabbit Monoclonal Antibody (OLIG2/7366R). Confirmation of Purity and Integrity of Antibody.



IHC analysis of FFPE human brain (RNA expression: 54.2 nTPM) stained with OLIG2 Recombinant Rabbit Monoclonal Antibody (OLIG2/7366R). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

Olig2, a basic helix-loop-helix transcription factor, is involved in oligodendroglial specification. Olig2 expression has been reported in most glial tumors, such as oligodendrogliomas and astrocytomas. Although more than half of glioblastomas are positive for Olig2, expression is very weak in terms of both percentage of labeled cells and intensity. No Olig2 expression has been found in the non-glial tumors including neuroepithelial tumors, ependymomas, subependymomas, medulloblastomas, and non-neuroepithelial tumors, such as CNS lymphomas, meningiomas, schwannomas, atypical teratoid/rhabdoid tumor, and haemangioblastomas. Compared to the strong staining seen in glioma samples, a weak expression is observed in non-tumoral brain tissue.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Neural Stem Cells
