

## Recombinant p57Kip2 (Mitotic Inhibitor/Suppressor Protein) Antibody

Mouse Monoclonal Antibody [Clone rKIP2/7238]

Catalog No	Format	Size
1028-MSM8-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1028-MSM8-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1028-MSM8-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


<b>Clone</b>	rKIP2/7238
<b>Gene Name</b>	CDKN1C
<b>Immunogen</b>	Recombinant full-length human p57Kip2 protein
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	57kDa
<b>Cellular Localization</b>	Nucleus.
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Human colon or prostate carcinomas. Brain.

\*Optimal dilution for a specific application should be determined.


### Product Images for Recombinant p57Kip2 (Mitotic Inhibitor/Suppressor Protein) Antibody




SDS-PAGE Analysis of Purified CDKN1C Recombinant Mouse Monoclonal Antibody(rKIP2/7238 ). Confirmation of Purity and Integrity of Antibody.



Western blot analysis of Human Brain tissue lysate using p57 Recombinant Mouse Monoclonal Antibody (rKIP2/7238).



Western blot analysis of Brain tissue lysates of different species using p57 Recombinant Mouse Monoclonal Antibody (rKIP2/7238).



IHC analysis of formalin-fixed, paraffin-embedded human bladder. Strong nuclear staining using rKIP2/7238 at 2ug/ml in PBS for 30min RT. Inset: PBS instead of primary antibody; secondary only negative control.

### Specificity & Comments

Recognizes a protein of 57kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab produced in HEK293 cell mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Endothelial Cell Marker, Infectious Disease, Lung Cancer, Nuclear Marker