

CD99 / MIC2 (Ewing s Sarcoma Marker) Antibody

Mouse Monoclonal Antibody [Clone MIC2/7866]

Catalog No	Format	Size
4267-MSM16-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4267-MSM16-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4267-MSM16-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	MIC2/7866
Gene Name	CD99
Immunogen	Recombinant fragment (around aa1-185) of human MIC2 protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	27-32kDa
Cellular Localization	Cell surface.
Species Reactivity	Human
Positive Control	MCF7, MOLT-4 cells. Human pancreas or Ewing s Sarcoma.

*Optimal dilution for a specific application should be determined.

Product Images for CD99 / MIC2 (Ewing s Sarcoma Marker) Antibody




Formalin-fixed, paraffin-embedded human pancreas stained with CD99 Mouse Monoclonal Antibody (MIC2/7866). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

SDS-PAGE Analysis of Purified CD99 Mouse Monoclonal Antibody (MIC2/7866). Confirmation of Purity and Integrity of Antibody.



Western blot analysis of MCF7 cell lysate using CD99 / MIC2 (Ewing sSarcoma Marker) Mouse Monoclonal Antibody (MIC2/7866).



Formalin-fixed, paraffin-embedded human pancreas stained with CD99 Mouse Monoclonal Antibody (MIC2/7866). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

Recognizes a sialoglycoprotein of 27-32kDa, identified as CD99, or MIC2 gene product, or E2 antigen. MIC2 gene is located in the pseudo-autosomal region of the human X and Y chromosome. MIC2 gene encodes two distinct proteins, which are produced by alternative splicing of the CD99 gene transcript and are identified as bands of 30 and 32kDa (p30/32). Although its function is not fully understood, CD99 is implicated in various cellular processes including homotypic aggregation of T cells, upregulation of T cell receptor and MHS molecules, apoptosis of immature thymocytes and leukocyte diapedesis. CD99 is expressed on the cell membrane of some lymphocytes, cortical thymocytes, and granulosa cells of the ovary. Most pancreatic islet cells, Sertoli cells of the testis, and some endothelial cells express this antigen. Mature granulocytes express very little or no CD99. MIC2 is strongly expressed on Ewing s sarcoma cells and primitive peripheral neuroectodermal tumors.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified by Protein A Column. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Immunology