

Recombinant SPARC / Osteonectin Antibody

Rabbit Monoclonal Antibody [Clone OSTN/8528R]

| Catalog No | Format | Size |
|------------------|--|--------|
| 6678-RBM17-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 6678-RBM17-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 6678-RBM17-P1ABX | Purified Ab WITHOUT BSA or Azide at 1.0mg/ml | 100 ug |



| Applications | Tested Dillution | Note |
|----------------------------|------------------|---|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details

| | |
|-------------------------------|---|
| Clone | OSTN/8528R |
| Immunogen | Recombinant fragment (around aa22-146) of the human Osteonectin protein (exact sequence is proprietary) |
| Host | Rabbit |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG / Kappa |
| Mol. Weight of Antigen | 34.63kDa |
| Cellular Localization | Basement membrane, Extracellular matrix, Extracellular space, Secreted |
| Species Reactivity | Human |
| Positive Control | Human placenta lymph node or spleen. JEG3 or A549 cell lysates. |

*Optimal dilution for a specific application should be determined.

Product Images for Recombinant SPARC / Osteonectin Antibody



SDS-PAGE Analysis of Purified Osteonectin Recombinant Rabbit Monoclonal Antibody (OSTN/8528R). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human tonsil stained with Osteonectin Recombinant Rabbit Monoclonal Antibody (OSTN/8528R). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

SPARC (for secreted protein acidic and rich in cysteine) is a phosphorylated, acidic, glycine-rich glycoprotein that is secreted by endothelial cells and is present in large amounts in the parietal endoderm of mouse embryos and in human placenta. It is identical to osteonectin, a protein important to bone calcification that is highly conserved between species. SPARC, which can be selectively expressed by the endothelium in response to certain types of injury, induces rounding in adherent endothelial cells in vitro. It regulates endothelial barrier function through F-Actin-dependent changes in cell shape, coincident with the appearance of intercellular gaps, which provide a paracellular pathway for extravasation of macromolecules.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
