

Recombinant STING1 / TMEM173 Antibody

Rabbit Monoclonal Antibody [Clone STING1/8129R]





Catalog No	Format	Size
340061-RBM12-P0	Purified Ab with BSA and Azide	200ug/ml
340061-RBM12-P1	Purified Ab with BSA and Azide	200ug/ml
340061-RBM12-P1ABX	Purified Ab WITHOUT BSA and Azide	1.0mg/ml



Applications	Tested Dillution
Immunohistochemistry (IHC)	1-2ug/ml

Product Details	
Clone	STING1/8129R
Gene Name	STING1
Immunogen	Recombinant fragment (around aa190-290) of human TMEM173 protein (exact sequence is proprietary)
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	37-50kDa
Cellular Localization	Cytoplasm.
Species Reactivity	Human
Positive Control	Cytoplasmic expression in several tissues including cells in respiratory tract fallopian tube and cells in lymphoid tissues.

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant STING1 / TMEM173 Antibody





Specificity & Comments

TMEM17 (transmembrane protein 173) is a 379 amino acid protein encoded by a gene mapping to human chromosome 5. With 181 million base pairs encoding around 1,000 genes, chromosome 5 is about 6% of human genomic DNA. It is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5 associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome. Deletion of 5q or chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

Research Areas

Immunology, Infectious Disease

Known Applications & Suggested Dilutions

Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT),(Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes),Optimal dilution for a specific application should be determined.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis.

There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C.Antibody is stable for 24 months. Non-hazardous. No MSDS required.