

## Recombinant CD45RA / LCA (B-Cell Marker) Antibody

Rabbit Monoclonal Antibody [Clone PTPRC/8699R]

Catalog No	Format	Size
5788-RBM52-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
5788-RBM52-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
5788-RBM52-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dilution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details


Clone	PTPRC/8699R
Gene Name	PTPRC
Immunogen	Recombinant full-length human PTPRC protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Mol. Weight of Antigen	180-220kDa
Cellular Localization	Cell Surface. Cytoplasm.
Species Reactivity	Human
Positive Control	Human tonsil or lymph node. Ramos

\*Optimal dilution for a specific application should be determined.


### Product Images for Recombinant CD45RA / LCA (B-Cell Marker) Antibody




Western Blot Analysis of Human Tonsil tissue lysate using CD45 Recombinant Rabbit Monoclonal Antibody (PTPRC/8699R).



SDS-PAGE Analysis of Purified Receptor-type tyrosine-protein phosphatase C Recombinant Rabbit Monoclonal Antibody (PTPRC/8699R). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of Ramos cell lysate using CD45 Recombinant Rabbit Monoclonal Antibody (PTPRC/8699R).



Formalin-fixed, paraffin-embedded human tonsil stained with CD45RA Recombinant Rabbit Monoclonal Antibody (PTPRC/8699R). Inset: PBS instead of primary antibody; secondary only negative control.

### Specificity & Comments

CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45R functions as a phosphor-tyrosine phosphatase. Antibody to CD45 is useful in differential diagnosis of lymphoid tumors from non-hematopoietic undifferentiated neoplasms.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Developmental Biology, Immunology, B Cell Markers, Dendritic Cell Marker, Hematopoietic Stem Cells, Mesenchymal Stem Cell Differentiation