

Myelin Basic Protein Antibody

Mouse Monoclonal Antibody [Clone MBP/4275]

Catalog No	Format	Size
4155-MSM5-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4155-MSM5-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4155-MSM5-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug

Applications	Tested Dillution	Note
Immunofluorescence (IF)	1-3ug/ml	
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	MBP/4275
Gene Name	MBP
Immunogen	Recombinant human MBP protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	13-21kDa
Cellular Localization	Myelin membrane; Peripheral membrane protein; Cytoplasmic side (of myelin).
Species Reactivity	Human
Positive Control	Human brain or astrocytoma tissue.

*Optimal dilution for a specific application should be determined.


Product Images for Myelin Basic Protein Antibody



Formalin-fixed, paraffin-embedded human brain stained with Myelin Basic Protein Mouse Monoclonal Antibody (MBP/4275).



SDS-PAGE Analysis of Purified Myelin Basic Protein Mouse Monoclonal Antibody (MBP/4275). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of human brain tissue lysate using Myelin Basic Protein Mouse Monoclonal Antibody (MBP/4275).



Formalin-fixed, paraffin-embedded human brain stained with Myelin Basic Protein Mouse Monoclonal Antibody (MBP/4275).

Specificity & Comments

Myelin basic protein (MBP) is the second most abundant protein in central nervous system (CNS) myelin: it comprises 30% of the total protein and about 10% of the dry weight of myelin. It is the only structural protein found so far to be essential for formation of CNS myelin, and has been called the executive molecule of myelin. MBP can interact with a number of polyanionic proteins including actin, tubulin, calmodulin, and clathrin, and negatively charged lipids, and acquires structure on binding to them. It may act as a membrane actin-binding protein, which might allow it to participate in transmission of extracellular signals to the cytoskeleton in oligodendrocytes and tight junctions in myelin. MBP may be applicable as a marker for oligodendrogliomas. MBP/4275 is used in clinical diagnostics to detect MBP levels or myelination in human MBP. MBP/4275 is used in a two-site ELISA with clone MBP/4273. Both are used interchangeably as capture and detection antibody.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Complement System, Developmental Biology, Neuroscience