

Recombinant MUC5AC (Mucin 5AC / Gastric Mucin) Antibody

Mouse Monoclonal Antibody [Clone rMUC5AC/8052]

Catalog No	Format	Size
4586-MSM17-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4586-MSM17-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4586-MSM17-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	rMUC5AC/8052
Gene Name	MUC5AC
Immunogen	Recombinant full-length human MUC5AC protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	400-600 kDa
Cellular Localization	Cytoplasm, Secreted
Species Reactivity	Human
Positive Control	MCF-7 cells. Human stomach.

**Optimal dilution for a specific application should be determined.*

Product Images for Recombinant MUC5AC (Mucin 5AC / Gastric Mucin) Antibody



SDS-PAGE Analysis of Purified Mucin-5AC Recombinant Mouse Monoclonal Antibody (rMUC5AC/8052). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human stomach stained with MUC5AC Recombinant Mouse Monoclonal (rMUC5AC/8052) at 2ug/ml. HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

Mucins are a group of high molecular weight glycoproteins consisting of a mucin core protein and O-linked carbohydrates. Mucin 6 carries GlcNAc β 1 \rightarrow 4Gal β 2R structures, indicating that β 1, 4-N-acetylglucosaminyltransferase is important to the formation of the mucous glycoproteins in vivo. Mucin 5AC is a gel-forming mucin that is secreted from surface mucous cells. Glucocorticoid is required for the expression of Mucin 5AC mRNA and high doses of hydrocortisone suppresses its expression. Additionally, asthmatic fluid stimulates Mucin 5AC synthesis several-fold. The pro-inflammatory cytokines IL-6 and TNF α stimulate Mucin 5AC secretion and thus contribute to the upregulation of mucin by chronic inflammation. Expression of Mucin 5AC is retinoic acid (RA)- or retinol-dependent, and RA control of mucin genes is mediated by the retinoid acid receptor RAR α and, to a lesser extent, by RAR β . Thyroid hormone binding to thyroid receptors inhibits Mucin 5AC gene expression. Mucin 5AC is also expressed in normal endocervical epithelium, small intestine, gastric cells (Lewis type 1) and gastric metaplasia and it is one of the major mucins in the ethmoid mucosa.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Immunology, Infectious Disease