

## Recombinant NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody

Mouse Monoclonal Antibody [Clone rNGFR/8824]

Catalog No	Format	Size
4804-MSM13-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
4804-MSM13-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
4804-MSM13-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

### Product Details

<b>Clone</b>	rNGFR/8824
<b>Gene Name</b>	NGFR
<b>Immunogen</b>	Recombinant full-length human p75 NGFR protein
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG1 / Kappa
<b>Mol. Weight of Antigen</b>	75kDa
<b>Cellular Localization</b>	Cell Surface. Cytoplasm.
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	and perineural cells of peripheral nerves or tumors of nerve sheath differentiation. Human melanomas. Schwann cells U87 cells. Neuronal axons

\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant NGF-Receptor (p75) / CD271 (Soft Tissue Tumor Marker) Antibody



SDS-PAGE Analysis of Purified NGFR Mouse Recombinant Monoclonal Antibody (rNGFR/8824). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human melanoma stained with NGFR Mouse Recombinant Monoclonal Antibody (rNGFR/8824). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

**Specificity & Comments**

It recognizes a glycoprotein of 75kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoid follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts.

---

**Limitations and Warranty**

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

---

**Supplied As**

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

---

**Storage and Stability**

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

---

**Research Areas**

AKT Signaling, Mesenchymal Stem Cell Differentiation, Neuroscience, Nuclear Marker, Signal Transduction, Stem Cell Differentiation

---