

## CREB3L4 Antibody

Mouse Monoclonal Antibody [Clone PCRP-CREB3L4-1A3]



| Catalog No        | Format  | Size   |
|-------------------|---|--------|
| 148327-MSM1-P0    | Purified Ab with BSA and Azide at 200ug/ml    | 20 ug  |
| 148327-MSM1-P1    | Purified Ab with BSA and Azide at 200ug/ml    | 100 ug |
| 148327-MSM1-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |

| Applications               | Tested Dillution    | Note  |
|----------------------------|---------------------|---|
| Flow Cytometry (Flow)      | 1-2ug/million cells |   |
| Immunofluorescence (IF)    | 1-3ug/ml            |   |
| Immunohistochemistry (IHC) | 1-2ug/ml            | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |

| Product Details        |   |
|------------------------|---|
| Clone                  | PCRP-CREB3L4-1A3  |
| Gene Name              | CREB3L4   |
| Immunogen              | Recombinant fragment (aa1-298) of human CREB3L4 protein |
| Host                   | Mouse   |
| Clonality              | Monoclonal  |
| Isotype / Light Chain  | IgG2c / Kappa   |
| Mol. Weight of Antigen | 44kDa   |
| Cellular Localization  | Nucleus.  |
| Species Reactivity     | Human   |
| Positive Control       | HeLa cells.   |


\*Optimal dilution for a specific application should be determined.

### Product Images for CREB3L4 Antibody



SDS-PAGE Analysis of Purified CREB3L4 Mouse Monoclonal Antibody (PCRP-CREB3L4-1A3). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human prostate cancer stained with CREB3L4 Mouse Monoclonal Antibody (PCRP-CREB3L4-1A3). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



Analysis of Protein Array containing more than 19,000 full-length human proteins using CREB3L4 Mouse Monoclonal (PCRP-CREB3L4-1A3). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to be specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.

Flow Cytometric Analysis of PFA-fixed HeLa cells. CREB3L4 Mouse Monoclonal Antibody (PCRP-CREB3L4-1A3) followed by goat anti-mouse IgG-CF488 (blue); unstained cells (red).

### Specificity & Comments

The ubiquitously expressed cellular upstream stimulatory factor (USF) consists of USF-1 and USF-2 polypeptides which independently exhibit site-specific DNA binding and are members of the c-Myc-related family of regulatory factors containing helix-loop-helix domains. USF also contains a leucine repeat that is required for efficient DNA binding. USF was originally identified as an upstream stimulatory factor that binds the core sequence CACGTG in the adenovirus late promoter. These findings, together with the demonstration of cooperative interaction between USF and the initiator binding protein, TFII-I, raises the possibility of a more general involvement of USF in transcriptional regulation. While expression of both USF-1 and USF-2 species is ubiquitous, different ratios of USF homo- and heterodimers are found in different cell types.

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

AKT Signaling, Nuclear Marker