

BCOR Antibody

Mouse Monoclonal Antibody [Clone BCOR/2372]

Catalog No	Format	Size
54880-MSM2-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
54880-MSM2-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
54880-MSM2-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details


Clone	BCOR/2372
Gene Name	BCOR
Immunogen	Recombinant fragment (around aa 100-400) of human BCOR protein (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	192kDa
Cellular Localization	Nucleus
Species Reactivity	Human
Positive Control	Ubiquitously expression.

*Optimal dilution for a specific application should be determined.


Product Images for BCOR Antibody



SDS-PAGE Analysis of Purified BCOR Mouse Monoclonal Antibody (BCOR/2372). Confirmation of Purity and Integrity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteins using BCOR Mouse Monoclonal Antibody (BCOR/2372). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Formalin-fixed, paraffin-embedded human tonsil stained with BCOR Mouse Monoclonal Antibody (BCOR/2372). Inset: PBS instead of primary antibody; secondary only negative control.

Specificity & Comments

Bcl-6, a transcriptional repressor, can promote or inhibit apoptosis depending on the cell type and also plays an important role in normal immune responses. Bcl-6 negatively regulates NF κ B expression, thereby inhibiting NF κ B-mediated cellular functions and is frequently found to be deregulated in non-Hodgkin's lymphoma. BCoR (Bcl-6 corepressor) is a 1,755 amino acid protein that associates with histone deacetylases (HDACs) to transcriptionally repress Bcl-6. With ubiquitous expression, BCoR is localized to the nucleus where it interacts with other proteins through its three ANK repeat domains. Mutations in the gene encoding BCoR result in microphthalmia with associated anomalies 2, also known as anophthalmia, which is characterized by variable features, such as renal aplasia, mental retardation, hyospadias, microencephaly and cryptorchidism. There are four isoforms of BCoR which are produced as a result of alternative splicing events.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.