

Recombinant mRNA cap guanine-N(7) methyltransferase Antibody

Rabbit Monoclonal Antibody [Clone TAG72/8317R]

Catalog No	Format	Size
182875-RBM11-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
182875-RBM11-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
182875-RBM11-P1ABX	Purified Ab WITHOUT BSA or Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	TAG72/8317R
Immunogen	Recombinant full-length human TAG-72 protein
Host	Rabbit
Clonality	Monoclonal
Isotype / Light Chain	IgG / Kappa
Species Reactivity	Human
Positive Control	Jurkat cells. Human breast or lung carcinoma.

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant mRNA cap guanine-N(7) methyltransferase Antibody



Formalin-fixed, paraffin-embedded human colon carcinoma stained with TAG-72 Rabbit Recombinant Monoclonal Antibody (TAG72/8317R). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.



Western blot analysis of Jurkat cell lysate using TAG-72 Recombinant Rabbit Monoclonal Antibody (TAG72/8317R).



SDS-PAGE Analysis of Purified TAG-72 Rabbit Recombinant Monoclonal Antibody (TAG72/8317R). Confirmation of Purity and Integrity of Antibody.

Specificity & Comments

Recognizes an oncofetal antigen of 220kDa, identified as a tumor-associated glycoprotein (TAG-72) with properties of a mucin. This MAb defines the mucin-carried sialylated-Tn epitope. TAG-72 is usually expressed by adenocarcinomas, but is negative in mesotheliomas. Studies have reported that this antibody has 80% sensitivity and 93% specificity for pulmonary adenocarcinoma. Therefore, TAG-72 is a useful marker to distinguish between mesothelioma and adenocarcinoma. However, false positive reactions can occur so results must be interpreted with the utmost caution. This antibody may be useful in the differentiation of non-small cell carcinomas from small cell carcinomas of the lung. The combined use of anti-TAG-72 and anti-GCDFP-15 is valuable in the diagnosis of apocrine carcinoma

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab produced in a mammalian-based expression system. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8 °C. Antibody without azide - store at -20 to -80 °C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.