

## Recombinant Creatine Phosphokinase-BB (CK-BB) Antibody

Rabbit Monoclonal Antibody [Clone CKBB/8310R]

Catalog No	Format	Size
1152-RBM15-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
1152-RBM15-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
1152-RBM15-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

### Product Details

<b>Clone</b>	CKBB/8310R
<b>Gene Name</b>	CKB
<b>Immunogen</b>	Recombinant human full-length CKB protein
<b>Host</b>	Rabbit
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG / Kappa
<b>Mol. Weight of Antigen</b>	42kDa
<b>Cellular Localization</b>	Cytoplasm.
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	Y79 or HEK293 cells. Human cerebellum.


\*Optimal dilution for a specific application should be determined.

### Product Images for Recombinant Creatine Phosphokinase-BB (CK-BB) Antibody




SDS-PAGE Analysis of Purified CKBB Recombinant Rabbit Monoclonal Antibody (CKBB/8310R). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human brain stained with CKBB Recombinant Rabbit Monoclonal Antibody (CKBB/8310R). Inset: PBS instead of primary antibody; secondary only negative control.



Western blot analysis of Y79 cell lysate using CKBB Rabbit Recombinant Monoclonal Antibody (CKBB/8310R).



Western Blot Analysis of HEK293 lysate using CKBB Recombinant Rabbit Monoclonal Antibody (CKBB/8310R ).

### Specificity & Comments

Creatine kinases (CK) are a large family of isoenzymes that regulate levels of ATP in subcellular compartments, where they provide ATP at sites of fluctuating energy demand by the transfer of phosphates between creatine and adenine nucleotides. CKs provide the energy of phosphate hydrolysis necessary to drive the normal function of many cellular systems. In cells, the cytosolic CK enzymes consist of two subunits, which can be either B (brain type) or M (muscle type). There are three different isoenzymes: CKMM, CKBB and CKMB. This MAb recognizes the CKBB isoenzyme and does not react with the B subunit in CKMB.

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cardiovascular, Signal Transduction

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.