

Coagulation Factor VII / F7 Antibody

Mouse Monoclonal Antibody [Clone F7/4931]

Catalog No	Format	Size
2155-MSM11-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2155-MSM11-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2155-MSM11-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details


Clone	F7/4931
Gene Name	F7
Immunogen	Recombinant fragment (around aa 366-466) of human F7 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2 / Kappa
Mol. Weight of Antigen	200/80/73/50/43kDa
Cellular Localization	Secreted.
Species Reactivity	Human
Positive Control	Human liver or gastric carcinoma.

*Optimal dilution for a specific application should be determined.


Product Images for Coagulation Factor VII / F7 Antibody



Western blot analysis of T47D cell lysate using Coagulation Factor VII Mouse Monoclonal Antibody (F7/4931).



Formalin-fixed, paraffin-embedded human kidney stained with Coagulation Factor VII Mouse Monoclonal Antibody (F7/4931). Inset: PBS instead of primary antibody; secondary only negative control.



SDS-PAGE Analysis Purified Coagulation Factor VII Mouse Monoclonal Antibody (F7/4931). Confirmation of Integrity and Purity of Antibody.

Specificity & Comments

Hemostasis following tissue injury involves the deployment of essential plasma procoagulants (prothrombin and Factors X, IX, V and VIII), which are involved in a blood coagulation cascade that leads to the formation of insoluble Fibrin clots and the promotion of platelet aggregation. Coagulation Factor VII (serum prothrombin conversion accelerator, proconvertin, F7, Factor VII) is a 406 amino acid, vitamin K-dependent, single chain serine protease that is synthesized in the liver and circulates as an inactive precursor. Factor IX A, Factor X A, Factor XII A or Thrombin-mediated proteolytic cleavage of Factor VII at Arg 152-Ile 153 generates Factor VII A, an active serine protease composed of a catalytic heavy chain disulfide linked to a light chain, containing two EGF-like domains. Mutations at the F7 locus that lead to Factor VII deficiencies are generally asymptomatic or phenotypically uncharacterized, with hemorrhagic diathesis occurring at extremely low levels.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Mitochondria Marker