

GAD2 / GAD65 (GABAergic Neuronal Marker) Antibody

Mouse Monoclonal Antibody [Clone GAD2/6484]

Catalog No	Format	Size
2572-MSM4-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
2572-MSM4-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
2572-MSM4-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug



Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes
Western Blot (WB)	2-4ug/ml	

Product Details

Clone	GAD2/6484
Gene Name	GAD2
Immunogen	Recombinant fragment of human GAD2 (GAD65) protein (around aa 1-200) (exact sequence is proprietary)
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG1 / Kappa
Mol. Weight of Antigen	65kDa
Cellular Localization	Cytoplasm.
Species Reactivity	Human
Positive Control	Human pancreas or brain.

*Optimal dilution for a specific application should be determined.

Product Images for GAD2 / GAD65 (GABAergic Neuronal Marker) Antibody



SDS-PAGE Analysis of Purified GAD2 (GAD65) Mouse Monoclonal Antibody (GAD2/6484). Confirmation of Purity and Integrity of Antibody.

Formalin-fixed, paraffin-embedded human thyroid stained with GAD2 (GAD65) Mouse Monoclonal Antibody (GAD2/6484). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

This MAb recognizes a protein of 65kDa, which is identified as glutamic acid decarboxylase 2 (GDA2). It is responsible for catalyzing the production of gamma-aminobutyric acid from L-glutamic acid. There are two forms of glutamic acid decarboxylases (GADs) that are found in the brain: GAD2 (also known as GAD65) and GAD1 (also known as GAD67). GAD1 and GAD2 are members of the group II decarboxylase family of proteins and are responsible for catalyzing the rate-limiting step in the production of GABA (?-aminobutyric acid) from L-glutamic acid. Although both GADs are found in the brain, GAD2 localizes to synaptic vesicle membranes in nerve terminals, while GAD1 is distributed throughout the cell. A pathogenic role for GAD2 is identified in the human pancreas since it has been identified as an autoantibody and an auto-reactive T cell target in insulin-dependent diabetes.?

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Neural Stem Cells, Neuroscience, Transcription Factors