

Annexin A2 Antibody

Mouse Monoclonal Antibody [Clone 3D5/4]

Catalog No	Format	Size
302-MSM1-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
302-MSM1-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
302-MSM1-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Immunohistochemistry (IHC)	1-2ug/ml	30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes

Product Details

Clone	3D5/4
Gene Name	ANXA2
Immunogen	Recombinant full-length human Annexin A2 protein
Host	Mouse
Clonality	Monoclonal
Isotype / Light Chain	IgG2a / Kappa
Mol. Weight of Antigen	36kDa
Cellular Localization	Secreted > extracellular space > extracellular matrix > basement membrane. Melosome. In the lamina beneath the plasma membrane. Identified by mass spectrometry in melosome fractions from
Species Reactivity	Human
Positive Control	HeLa or A549 cells. Human colon cancer.

*Optimal dilution for a specific application should be determined.

Product Images for Annexin A2 Antibody



Formalin-fixed, paraffin-embedded human colon cancer stained with Annexin A2 Mouse Monoclonal Antibody (3D5/4). HIER: Tris/EDTA, pH9.0, 45min. 2°C: HRP-polymer, 30min. DAB, 5min.

Specificity & Comments

The Annexin family of calcium-binding proteins is composed of at least ten mammalian genes and is characterized by a conserved core domain which binds phospholipids in a Ca²⁺-dependent manner and a unique amino-terminal region which may confer binding specificity. The interaction between these proteins and biological membranes has led to the hypothesis that they are involved in cellular trafficking processes such as endocytosis, exocytosis and cellular adhesion. Annexin I, alternatively referred to as lipocortin, has been implicated as a mediator of the anti-inflammatory response produced by glucocorticoids and as an inhibitor of cPLA₂, a potent mediator of inflammation. Annexin II, also called p36, exists as a monomer or as a heterotetramer, complexed with the S-100-related protein p11. This complex is termed Calpactin I. In the tetrameric form, Annexin II is an efficient substrate of PKC family and Src pp60.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

Cardiovascular, Cytokine Signaling, Immunology