

Recombinant HLA-DR (MHC II) Antibody

Rabbit Monoclonal Antibody [Clone HLA-DRA/8287R]

| Catalog No | Format | Size |
|------------------|-----------------------------------------------|--------|
| 3122-RBM10-P0 | Purified Ab with BSA and Azide at 200ug/ml | 20 ug |
| 3122-RBM10-P1 | Purified Ab with BSA and Azide at 200ug/ml | 100 ug |
| 3122-RBM10-P1ABX | Purified Ab WITHOUT BSA and Azide at 1.0mg/ml | 100 ug |


| Applications | Tested Dillution | Note |
|----------------------------|------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Immunohistochemistry (IHC) | 1-2ug/ml | 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes |
| Western Blot (WB) | 2-4ug/ml | |

Product Details


| | |
|-------------------------------|-----------------------------------------------------------------------------------------------|
| Clone | HLA-DRA/8287R |
| Gene Name | HLA-DRA |
| Immunogen | Recombinant fragment (around aa1-200) of human HLA-DR protein (exact sequence is proprietary) |
| Host | Rabbit |
| Clonality | Monoclonal |
| Isotype / Light Chain | IgG / Kappa |
| Mol. Weight of Antigen | ~28kDa (beta chain) |
| Cellular Localization | Cell surface. |
| Species Reactivity | Human |
| Positive Control | Human tonsil or lymph node. |

*Optimal dilution for a specific application should be determined.


Product Images for Recombinant HLA-DR (MHC II) Antibody



SDS-PAGE Analysis of Purified HLA-DR Recombinant Rabbit Monoclonal(HLA-DRA/8287R). Confirmation of Purity and Integrity of Antibody.



Western Blot Analysis of human tonsil lysate using HLA-DRA Recombinant Rabbit Monoclonal Antibody (HLA-DRA/8287R).



Formalin-fixed, paraffin-embedded human tonsil stained with HLA-DR Recombinant Rabbit Monoclonal Antibody (HLA-DRA/8287R). Inset: PBS instead of primary antibody; secondary only negative control

Specificity & Comments

This MAbs reacts with the beta-chain of HLA-DR antigen, a member of MHC class II molecules. It does not cross react with HLA-DP and HLA-DQ. The L243 antibody recognizes a different epitope than the HLA-DRA/8287R monoclonal antibody, and these antibodies do not cross-block binding to each other's respective epitopes. HLA-DR is a heterodimeric cell surface glycoprotein comprised of a 36kDa alpha (heavy) chain and a 28kDa beta (light) chain. It is expressed on B-cells, activated T-cells, monocytes/macrophages, dendritic cells and other non-professional APCs. In conjunction with the CD3/TCR complex and CD4 molecules, HLA-DR is critical for efficient peptide presentation to CD4+ T cells. It is an excellent histiocytic marker in paraffin sections producing intense staining. True histiocytic neoplasms are similarly positive. HLA-DR antigens also occur on a variety of epithelial cells and their corresponding neoplastic counterparts.

Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.

Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

Research Areas

B Cell Markers, Cardiovascular, Cytokine Signaling, Dendritic Cell Marker, Hematopoietic Stem Cells, Immunology