

## BCL6 Antibody

Mouse Monoclonal Antibody [Clone PCR-P-BCL6-1B1]

Catalog No	Format	Size
604-MSM17-P0	Purified Ab with BSA and Azide at 200ug/ml	20 ug
604-MSM17-P1	Purified Ab with BSA and Azide at 200ug/ml	100 ug
604-MSM17-P1ABX	Purified Ab WITHOUT BSA and Azide at 1.0mg/ml	100 ug


Applications	Tested Dillution	Note
Flow Cytometry (Flow)	1-2ug/million cells	

### Product Details

<b>Clone</b>	PCR-P-BCL6-1B1
<b>Gene Name</b>	BCL6
<b>Immunogen</b>	Recombinant fragment (around aa528-601) of human BCL6 protein
<b>Host</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Isotype / Light Chain</b>	IgG2b
<b>Mol. Weight of Antigen</b>	95kDa
<b>Cellular Localization</b>	Nucleus. Mitochondria.
<b>Species Reactivity</b>	Human
<b>Positive Control</b>	HeLa or MCF-7 cells.

\*Optimal dilution for a specific application should be determined.


### Product Images for BCL6 Antibody



SDS-PAGE Analysis. Purified BCL6 Mouse Monoclonal Antibody (PCR-P-BCL6-1B1). Confirmation of Integrity and Purity of Antibody.



Analysis of Protein Array containing more than 19,000 full-length human proteins using BCL6 Mouse Monoclonal Antibody (PCR-P-BCL6-1B1). Z- and S- Score: The Z-score represents the strength of a signal that a monoclonal antibody (MAb) (in combination with a fluorescently-tagged anti-IgG secondary antibody) produces when binding to a particular protein on the HuProt™ array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If targets on HuProt™ are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-score. S-score therefore represents the relative target specificity of a MAb to its intended target. A MAb is considered to specific to its intended target, if the MAb has an S-score of at least 2.5. For example, if a MAb binds to protein X with a Z-score of 43 and to protein Y with a Z-score of 14, then the S-score for the binding of that MAb to protein X is equal to 29.



Flow Cytometric Analysis of PFA-fixed HeLa cells. BCL6 Mouse Monoclonal Antibody (PCRP-BCL6-1B1) followed by goat anti-mouse IgG-CF488 (blue); unstained cells (red).

### Specificity & Comments

Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large (L&H) cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.

### Supplied As

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

### Storage and Stability

Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.

### Research Areas

Cardiovascular, Cytokine Signaling, Immunology, Nuclear Marker, Transcription Factors

### Limitations and Warranty

This antibody is available for research use only and is not approved for use in diagnosis. There are no warranties, expressed or implied, which extend beyond this description. Company is not liable for any personal injury or economic loss resulting from this product.